



Dietary protein and muscle in older persons

Douglas Paddon-Jones^a and Heather Leidy^b

Purpose of review

The purpose of this study is to highlight recent advances in nutrition and protein research that have the potential to improve health outcomes and status in ageing adults.

Recent findings

The beneficial effects of dietary protein on muscle health in older adults continue to be refined. Recent research has bolstered support for moderately increasing protein consumption beyond the current Recommended Dietary Allowance by adopting a meal-based approach in lieu of a less specific daily recommendation. Results from muscle protein anabolism, appetite regulation and satiety research support the contention that meeting a protein threshold (approximately 30 g/meal) represents a promising strategy for middle-aged and older adults concerned with maintaining muscle mass while controlling body fat.

Summary

Optimizing dietary protein intake to improve health requires a detailed consideration of topics including muscle protein anabolism, appetite control and satiety. Although each area of research continues to advance independently, recent collaborative and translational efforts have highlighted broad, translational consistencies related to the daily distribution and quantity of dietary protein.

Keywords

ageing, nutrition, sarcopenia, satiety, strength

OVERVIEW

Higher protein diets continue to gain scientific support as a strategy to preserve lean mass, promote weight loss, prevent weight-regain following weight loss or to simply maintain a healthy weight throughout the lifespan. The effectiveness of these diets appears to be driven, in part, by improvements in protein anabolism, daily appetite control and satiety. This brief synopsis will first summarize the state of the research regarding the effects of increased protein consumption on changes in body weight/composition during controlled-feeding trials and in free-living older adult populations. Recent conceptual and mechanistic advances in protein synthesis, energy expenditure and the regulation of food intake will also be explored. Finally, the impact of protein quantity and timing of consumption will be addressed.

SARCOPENIA AND PROTEIN RECOMMENDATIONS

A reduction in muscle mass and functional capacity is a largely inevitable consequence of ageing. The uncomplicated progression of sarcopenia results in a 3–8% reduction in muscle mass per decade, starting in the fourth or fifth decade of life. During this early

period, small decrements in muscle mass or function may be readily masked by subtle lifestyle adaptations. However, advanced sarcopenia is synonymous with physical frailty and is associated with an increased risk of falls and impairment in the ability to perform routine activities of daily living [1,2].

The Recommended Dietary Allowance (RDA) for protein (i.e. 0.8 g protein/kg/day) describes the quantity of protein required daily to prevent deficiency for all adults, irrespective of age. In recent years, there has been increased support for the contention that the current RDA for protein is insufficient to promote optimal health. To this end, several recent reviews and consensus statements have suggested that a protein intake between 1.0 and 1.5 g/kg/day may confer health benefits beyond

^aDepartment of Nutrition and Metabolism, University of Texas Medical Branch, Galveston, Texas and ^bDepartment of Nutrition & Exercise Physiology, University of Missouri, Columbia, Missouri, USA

Correspondence to Douglas Paddon-Jones, PhD, Department of Nutrition and Metabolism, The University of Texas Medical Branch, 301 University Blvd, Galveston, TX 77555-1124, USA. Tel: +1 409 772 3073; e-mail: djpaddon@utmb.edu

Curr Opin Clin Nutr Metab Care 2014, 17:5–11

DOI:10.1097/MCO.000000000000011

KEY POINTS

- Sarcopenia has debilitating consequences for the elderly community and considerable impact on healthcare systems.
- Moderate protein diets for young and older adults have led to greater weight loss, fat loss and preservation of lean mass compared with standard protein diets.
- Potential protein-related mechanism(s)-of-action include the increase in protein anabolism, appetite regulation and satiety.
- Meeting a protein threshold (approximately 30 g/meal) represents a promising strategy or dietary framework for middle-aged and older adults concerned with maintaining muscle mass while controlling body fat.
- Multidisciplinary research efforts are needed to provide practical, translatable data on the regulation of body composition and maintenance of muscle mass and function during ageing.

those afforded by simply meeting the minimum [3–8].

Analysis of large data sets such as the National Health and Nutrition Examination Survey (NHANES) suggests that many adult populations already consume a moderate-to-high protein diet (i.e. 1.3 g protein/kg/day) [9,10]. Unfortunately, older adults generally consume less protein and energy than their younger peers [6]. Approximately one-third of adults over 50 years of age fail to meet the RDA for protein, while approximately 10% of older women fail to meet even the Estimated Average Requirement (EAR) for protein (0.66 g protein/kg/day) [3,5]. For a 65 kg adult, the EAR represents a paltry 40 g of protein/day. Increasing and optimizing protein intake may be particularly important for older populations experiencing catabolic stressors such as illness, physical inactivity or injury [3,6,11,12]. In circumstances in which the capacity or ability to exercise is limited, nutrition, and protein consumption in particular, represents one of the few remaining options to improve muscle protein anabolism and ultimately preserve muscle mass and function.

Although there are compelling arguments supporting the role of higher protein diets for muscle health, it is equally clear that a growing proportion of the population chronically exceed their daily energy requirements [10,13]. In older adults, the outwardly paradoxical increase in the incidence of sarcopenic obesity highlights the need to look beyond the quantity of protein and energy consumed daily [14–16]. Recent research efforts

have clearly established the benefits of increased dietary protein within an energy-controlled diet for weight management [14,17]. However, relatively few studies have examined issues such as protein distribution [18,19,20], or adopted an integrated approach to examine the concomitant effects of increased protein on a broad, cross-disciplinary array of outcomes/themes such as protein metabolism, cell signalling, body composition, satiety, glucose regulation and overall macronutrient consumption.

BODY WEIGHT/COMPOSITION

Three recent meta-analyses examined the effects of higher protein diets on changes in body weight and body composition. Wycherley *et al.* [21] analysed 24 randomized controlled trials that compared higher protein or standard protein energy-restriction diets of 12.1 ± 9.3 weeks in duration in 1063 overweight and obese individuals between 18 and 80 years of age. The higher protein diets contained between 27 and 35% of daily energy intake as protein (1.07–1.60 g protein/kg/day), whereas the standard protein diets contained 16–21% protein (0.55–0.88 g protein/kg/day) [21]. Across this broad age spectrum, the higher protein diets led to greater weight loss [mean difference: -0.79 kg; 95% confidence interval (CI) -1.50 to -0.08 ; $P < 0.03$] and fat loss (mean difference: -0.87 kg; 95% CI -1.26 to -0.48 ; $P < 0.001$) than the standard protein diets [21]. Of particular importance for middle-aged and older adults, the higher protein diets were also associated with greater lean mass retention (during energy restriction) (mean difference: $+0.43$ kg; 95% CI 0.09 – 0.78 ; $P < 0.01$) than the standard protein diets [21].

Similar findings were also reported in a meta-analysis of studies involving 418 middle-aged adults (46–63 years) with type 2 diabetes [22]. In this analysis, the higher protein diets, containing 25–32% protein, led to greater weight loss (mean difference: -2.08 kg; 95% CI -3.25 to -0.90) than diets containing 15–20% protein [22]. Santesso *et al.* [23] extended these findings to encompass 74 randomized controlled trials comparing higher protein (16–45%) vs. standard protein diets (5–23%). Irrespective of factors such as total energy consumption, health status and age, higher protein diets facilitated greater weight loss (mean difference: -0.36 kg; 95% CI -0.56 to -0.17 ; $P < 0.001$), reductions in BMI (mean difference: -0.37 kg/m²; 95% CI -0.56 to -0.19 ; $P < 0.001$) and reductions in waist circumference (mean difference: -0.43 cm; 95% CI -0.69 to -0.16 ; $P < 0.001$) compared with the standard protein diets. The magnitude of

change, although modest, has clear clinical relevance in light of the ongoing obesity epidemic as well as the increased prevalence of type 2 diabetes, metabolic syndrome, sarcopenia and sarcopenic obesity.

MUSCLE LOSS AND INACTIVITY: IS 50 THE NEW 70?

Establishing healthy exercise and nutrition behaviours should not be viewed as a switch that can be turned on at a particular age or in response to a health scare. Age-related conditions such as osteoporosis and sarcopenia do not develop acutely. Rather, they are insidious conditions whose onset may be accelerated by less than optimal lifestyle practices in early-middle age [24]. Nevertheless, middle-aged adults (i.e. 40–65 years) are comparatively underrepresented in research, falling between the typical age-based group assignments of ‘young’ (20–40 years) or ‘elderly’ (more than 65 years).

In a compilation of studies examining the effects of physical inactivity/bed rest on skeletal muscle, we nominally compared the rate and magnitude of muscle loss in young [25] and older adults [26] with a recently completed study in middle-aged adults [27,28] (Table 1).

Although controlled experimental conditions are certainly needed to confirm these data, it appears that, although healthy middle-age adults often have ‘normal’ or ‘youthful’ physiological responses during acute muscle metabolism studies [29,30], a metabolic perturbation or catabolic insult (e.g. inactivity, injury, malnutrition) may facilitate anabolic resistance or an ageing phenotype, increasing the rate of muscle loss.

These observations are consistent with the ‘Catabolic Crisis’ model of muscle loss we recently put forward [13] and suggest that prevention and treatment strategies targeting older adults could be extended to middle-aged adults who may also be at an increased risk of accelerated muscle loss during periods of physiological stress [31,32].

REGULATION OF PROTEIN INTAKE

Ingestive behaviour is a complex system composed of both physiological eating and reward-driven (i.e.

hedonic) eating. Physiological eating occurs in response to acute and/or chronic fluctuations in energy balance during times of fasting, meal omission and weight loss, whereas hedonic eating typically occurs in response to external (environmental) signals that stimulate memories and thoughts of food for enjoyment and/or reward. In the current, food-centric obesogenic environment, many support the premise that hedonic eating is the most significant factor in terms of overweight and weight gain; however, during weight loss (or following weight loss), the physiological signals that control hunger and satiety play a pivotal role. Thus, it is critical to identify the effects of dietary interventions on both types of eating. Recent evidence suggests that increased dietary protein modulates both systems.

Results from the majority of acute single-meal studies illustrate significant postprandial reductions in perceived hunger and increased perceived fullness following the consumption of higher vs. standard protein meals [33]. These responses are accompanied by hormonal responses including reductions in the hunger-stimulating hormone ghrelin and increases in the satiety-stimulating hormones peptide YY (PYY) and glucagon-like peptide-1 (GLP-1) [33,34]. Although these findings are more physiological in nature, recent data from our laboratory also illustrate that the consumption of higher protein meals significantly impacts reward-driven eating behaviour. Using functional MRI (fMRI), we have identified neural activation in response to rewarding food stimuli prior to subsequent eating occasions. Compared with standard protein meals, the higher protein versions lead to reduced activation in select cortico-limbic brain regions, including the insula, hippocampus, parahippocampus and/or middle prefrontal cortex, which control food motivation, reward and cravings as well as executive function [35,36]. Collectively, these data support the beneficial effect of dietary protein to modulate appetite control, satiety and food motivation/reward.

It is important to note that studies examining the regulation of food intake have been performed almost exclusively in young to middle-age men and women. Due to the well documented anorexic effect of ageing, the consumption of higher protein meals

Table 1. Comparison of leg lean mass loss per day during bed rest for young [25], middle-aged [27] and older adults [28]

	Bed rest (days)	Age (years)	Muscle loss (kg)	Rate of loss (g/day)
Young	28	38 ± 8	-0.40 ± 0.10	-14
Middle-age	14	52 ± 4	-1.16 ± 0.14	-83
Elderly	10	67 ± 5	-0.95 ± 0.15	-95

Table 2. Age-related appetite responses following standard vs. high protein meals [37]

	Age (years)	4-h postmeal appetite ^a following a standard protein meal (area under the curve; mm ² 240 min)	4-h postmeal appetite ^a following a high protein meal (area under the curve; mm ² 240 min)
Young	38 ± 2	7878 ± 1310	6071 ± 1145 ^b
Middle-aged	50 ± 1	6065 ± 1205	5432 ± 1053 ^b
Elderly	64 ± 2	5551 ± 1254 ^c	4646 ± 1096 ^{b,c}

^aAppetite was measured from 100 mm visual analogue scales assessing perceived hunger as 'how strong is your feeling of hunger', with end anchors of 'extremely' to 'not at all'. Questionnaires completed every 30 min throughout the 4-h period.

^bHigh protein vs. standard protein, $P < 0.05$.

^cOld vs. young, $P < 0.05$.

might exert an even larger reduction in appetite in the elderly. However, it is also possible that older individuals will be insensitive to increased protein consumption due to their already blunted, 'nonresponsive' appetite signalling. A retrospective analysis comparing the postmeal appetite responses following standard protein vs. higher protein meals across the lifespan is summarized in Table 2 [37]. Although the older individuals experienced blunted appetite compared with their younger counterparts, the consumption of higher protein meals led to greater reductions in perceived appetite than a standard protein meal in all age groups (Table 2; all, $P < 0.05$).

However, irrespective of age, consumption of moderately higher protein meals still led to reduced postprandial appetite responses compared with standard protein versions. Although muscle protein metabolism research consistently demonstrates the benefits of a moderately higher dietary protein intake, they do not address complex behavioural issues such as the drive to eat. Studies examining the acute and chronic effects of increased dietary protein on physiological and hedonic eating are needed in the elderly to better understand the anorexia of ageing to establish specific, age-appropriate protein recommendations.

PROTEIN QUANTITY

One of the remaining overarching questions in the field of protein research centres on the quantity of protein required to optimize muscle protein anabolism, appetite control and body weight management. The difficulty in answering this question lies with the multiple ways in which protein consumption can be quantified. The dietary guideline recommendations are reported as grams of protein/kg of body weight per day with a minimal quantity of 0.8 g protein/kg/day established as the quantity to prevent deficiencies. The dietary guidelines also include an acceptable macronutrient distribution range (AMDR). The AMDR for protein represents

10–35% of an individual's total daily energy intake and is defined as 'the range of intake for a particular energy source (i.e. protein) that is associated with reduced risk of chronic disease while providing intakes of essential nutrients'. Unfortunately, the prescriptive usefulness of the AMDR is diminished by the overly restrictive and generous quantities of protein represented by both ends of the range. Nevertheless, the protein meta-analyses presented in this synopsis suggest that the quantity of protein necessary to promote improvements in body composition falls between 1.2 and 1.6 g protein/kg/day (i.e. ~25–30% of daily intake as protein) [38]. This is well within the acceptable macronutrient range for protein and allows for the ability to meet the recommendations for other requirements including fruits, vegetables, dairy and fibre.

Many mechanistic studies and clinical trials examining protein synthesis, appetite control and satiety focus on meal-specific protein consumption and not on total daily intake. For some outcome measures, this may be a critical point. For example, the human body has limited capacity to temporarily store excess protein from a single large meal and use it to acutely stimulate muscle protein anabolism at a later date. In several recent studies and reviews, there is general agreement that approximately 30 g of protein/eating occasion is required to elicit an optimal or measureable change in a number of outcome variables [6,39–42]. For example, Fig. 1 shows the 2 h postprandial fullness response following the consumption of 350 kcal meals containing varying quantities of dietary protein (unpublished data, composite from previous studies with similar populations and experimental designs).

Although the consumption of all meals led to an immediate rise in fullness, the meal containing 30 g of protein elicited a larger (and more sustained) increase over the 2 h postprandial period compared with the other quantities. Thus, these data support a within-meal protein threshold to elicit increases in satiety. If three to four meals containing 30 g of protein/meal are consumed throughout the day,




FIGURE 1. Postprandial fullness responses following 350kcal meals varying in protein content. AUC, area under the curve ($\text{mm} \times 120 \text{ min} - 10$).

the total amount of protein equates to the quantities shown to elicit body weight/body composition changes described above [20]. Ingestion of larger protein meals may be justified for adults with increased energy demands or high protein turnover (e.g. athletes, some patient populations), but any potential benefit should be weighed against the risk of exceeding daily energy requirements.

TIMING AND DISTRIBUTION OF CONSUMPTION

Current diet trends continue to focus on the concept of eating frequency to promote optimal weight loss/body composition changes. Ironically, these trends cross the gamut of eating frequency with some advocating small 'mini' meals spread through the day, while others emphasize intermittent fasting, which includes meal omission, particularly breakfast. However, on the basis of the previous discussion, the quantity and distribution of protein embedded within this eating pattern may significantly impact the outcomes.

In older adults, two recent studies have focused on changes in muscle mass associated with protein

distribution patterns [18[■],19]. Bollwein *et al.* [18[■]] performed a cross-sectional analysis of groups of frail, prefrail and nonfrail adults over the age of 75 years. They noted that although total daily protein consumption was similar in all groups, nonfrail older adults consumed an evenly distributed protein diet (i.e. moderate amounts of protein at each meal). In contrast, frail and prefrail cohorts followed a more skewed protein ingestion pattern, consuming the bulk of their total daily protein during the midday meal.

A recent clinical trial asked a similar question, but found quite different results. Bouillanne *et al.* [19] conducted a 6-week randomized trial in hospitalized, older adults (average 85 years). They reported that patients who were provided with a 'pulsed' protein diet (8 a.m.: 4.5 g, noon: 47.8 g, 4 p.m.: 2.3 g, 7 p.m.: 10.9 g) experienced a significant improvement in lean mass compared with the 'spread' protein diet (8 a.m.: 12.2 g, noon: 21 g, 4 p.m.: 13.5 g, 7 p.m.: 21.2 g) that provided the same total amount of protein each day (i.e. 1.31 g protein/kg/day). Methodological differences aside, it is difficult to immediately reconcile the apparent discrepancy in the results of these studies.

However, an obvious area for further investigation centres on the threshold amount of protein per meal required by various ageing populations (healthy and clinical) to elicit a meaningful increase in muscle protein anabolism. In the case of the Bouillanne study, one could suggest that each of the spread protein meals (12–21 g protein) failed to meet the threshold required to optimally stimulate muscle protein anabolism [6,39–42], whereas the noon meal in the pulsed protein likely had more than enough protein (~48 g) to elicit a single, robust anabolic response [39].

We previously examined whether six small meals (i.e. 350 kcal/meal) consumed every 2 h improve daily appetite and satiety responses compared with the standard three larger meals (i.e. 700 kcal/meal) [43,44]. Protein quantity was also varied within each meal pattern. Overall, we found that protein quantity had a more robust effect on satiety than eating frequency. Higher protein meals led to greater feelings of fullness and PYY concentrations throughout the day than the normal protein meals. These data refute claims that consuming 'mini' meals and/or frequent snacking throughout the day is beneficial in controlling appetite and satiety. Further, these data support the 'protein threshold' concept of at least 30 g protein to elicit satiety responses, as the meals containing less than 30 g protein, as was the case for the six small high protein meal and the three large normal protein meal patterns, led to the smallest satiety response.

A recent longer term study by Arciero *et al.* [45] extended these findings to examine the effects of increased protein intake and meal frequency on changes in body weight and body composition. In this study, the higher protein diet groups, regardless of eating frequency, led to greater fat loss, particularly abdominal fat, than the normal protein diet. The higher protein diets also led to increased lean mass and reduced daily hunger vs. normal protein versions. The protein meal pattern in this study contained approximately 30 g of protein per eating occasion.

At the other end of the eating frequency spectrum is the concept of meal omission/skipping. Although intermittent fasting, which is a type of reduced eating frequency, is gaining traction in the lay press, limited data exist to support this dietary pattern. On the contrary, meal omission, particularly the breakfast meal, has been strongly associated with increased BMI, unhealthy weight gain and obesity [46]. We recently found that the consumption of breakfast led to increased appetite control, satiety and reduced reward-driven eating behaviour throughout the day compared with skipping the

morning meal [38]. However, a protein-rich breakfast, containing 35 g of protein, led to further improvements through greater increases in satiety throughout the day along with greater reductions in food motivation/reward than a normal protein breakfast containing only 13 g protein. Further, only the higher protein breakfast reduced unhealthy, evening snacking on high-fat snacks by approximately 200 kcal compared with skipping breakfast or consuming a normal protein breakfast. These data suggest that substantial protein intake at the breakfast meal appears to have immediate and sustained effects throughout the day. Previous data from our laboratory further support these findings by illustrating that protein at the morning meal leads to greater immediate and sustained satiety throughout the day than protein at lunch or dinner [47].

CONCLUSION

Sarcopenia has debilitating consequences for the elderly community and considerable impact on healthcare systems. The benefits of moderate protein diets for young and older adults have been well established and many ongoing research efforts seek to refine and incrementally add to our understanding. However, it is increasingly clear that multi-disciplinary research efforts are needed to provide practical, translatable data on the regulation of body composition and maintenance of muscle mass and function during ageing. A compilation of data from recent muscle protein anabolism, appetite regulation and satiety studies suggest that meeting a protein threshold (approximately 30 g/meal) represents a promising strategy or dietary framework for middle-aged and older adults concerned with maintaining muscle mass while controlling body fat.

Acknowledgements

This research was supported by grants from the NIH/NINR RO1 NR012973 (D.P.J.), UTMB Claude D. Pepper Older Americans Independence Center #P30 AG024832 (D.P.J.), Pork Checkoff (H.L.), Egg Nutrition Center (H.L.) and Beef Checkoff (H.L.).

Conflicts of interest

D.P.J. has received compensation for speaking engagements with Abbott Nutrition, the National Cattlemen's Beef Association, the American Egg Board and the National Dairy Council. H.L. has received compensation for speaking engagements with the National Cattlemen's Beef Association, the American Egg Board and the National Dairy Council.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Landi F, Liperoti R, Russo A, *et al.* Sarcopenia as a risk factor for falls in elderly individuals: results from the iSIRENTE study. *Clin Nutr* 2012; 31:652–658.
 2. Fisher SR, Goodwin JS, Protas EJ, *et al.* Ambulatory activity of older adults hospitalized with acute medical illness. *J Am Geriatr Soc* 2011; 59:91–95.
 3. Wolfe RR, Miller SL. The recommended dietary allowance of protein: a misunderstood concept. *JAMA* 2008; 299:2891–2893.
 4. Wolfe RR, Miller SL, Miller KB. Optimal protein intake in the elderly. *Clin Nutr* 2008; 27:675–684.
 5. Houston DK, Nicklas BJ, Ding J, *et al.* Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *Am J Clin Nutr* 2008; 87:150–155.
 6. Volpi E, Campbell WW, Dwyer JT, *et al.* Is the optimal level of protein intake for older adults greater than the recommended dietary allowance? *J Gerontol A Biol Sci Med Sci* 2013; 68:677–681.
 7. Fielding RA, Vellas B, Evans WJ, *et al.* Sarcopenia: an undiagnosed condition in older adults. Current consensus definition: prevalence, etiology, and consequences. International working group on sarcopenia. *J Am Med Dir Assoc* 2011; 12:249–256.
 8. Phillips SM. Dietary protein requirements and adaptive advantages in athletes. *Br J Nutr* 2012; 108 (Suppl 2):S158–S167.
 9. Fulgoni VL. Current protein intake in America: analysis of the National Health and Nutrition Examination Survey, 2003–2004. *Am J Clin Nutr* 2008; 87:1554S–1557S.
 10. Layman DK. Dietary guidelines should reflect new understandings about adult protein needs. *Nutr Metab (Lond)* 2009; 6:12.
 11. English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Curr Opin Clin Nutr Metab Care* 2010; 13:34–39.
 12. Wall BT, Dirks ML, van Loon LJ. Skeletal muscle atrophy during short-term disuse: implications for age-related sarcopenia. *Ageing Res Rev* 2013; 12:898–906.
 13. Drewnowski A, Fulgoni VL, Young MK, Pitman S. Nutrient-rich foods: applying nutrient navigation systems to improve public health. *J Food Sci* 2008; 73:H222–H228.
 14. Tang M, Armstrong CL, Leidy HJ, Campbell WW. Normal vs. high-protein weight loss diets in men: effects on body composition and indices of metabolic syndrome. *Obesity (Silver Spring)* 2013; 21:E204–E210.
 15. Coker RH, Miller S, Schutzler S, *et al.* Whey protein and essential amino acids promote the reduction of adipose tissue and increased muscle protein synthesis during caloric restriction-induced weight loss in elderly, obese individuals. *Nutr J* 2012; 11:105.
 16. Fielding RA, Gunstad J, Gustafson DR, *et al.* The paradox of overnutrition in aging and cognition. *Ann N Y Acad Sci* 2013; 1287:31–43.
 17. Leidy HJ, Tang M, Armstrong CL, *et al.* The effects of consuming frequent, higher protein meals on appetite and satiety during weight loss in overweight/obese men. *Obesity (Silver Spring)* 2011; 19:818–824.
 18. Bollwein J, Diekmann R, Kaiser MJ, *et al.* Distribution but not amount of protein intake is associated with frailty: a cross-sectional investigation in the region of Nurnberg. *Nutr J* 2013; 12:109.
- This cross-sectional study highlights that even when older adults exceed the RDA for protein, an even protein distribution may be associated with muscle loss/frailty.
19. Bouillanne O, Curis E, Hamon-Vilcot B, *et al.* Impact of protein pulse feeding on lean mass in malnourished and at-risk hospitalized elderly patients: a randomized controlled trial. *Clin Nutr* 2013; 32:186–192.
 20. Paddon-Jones D, Rasmussen BB. Dietary protein recommendations and the prevention of sarcopenia. *Curr Opin Clin Nutr Metab Care* 2009; 12:86–90.
 21. Wycherley TP, Moran LJ, Clifton PM, *et al.* Effects of energy-restricted high-protein, low-fat compared with standard-protein, low-fat diets: a meta-analysis of randomized controlled trials. *Am J Clin Nutr* 2012; 96:1281–1298.
 22. Dong JY, Zhang ZL, Wang PY, Qin LQ. Effects of high-protein diets on body weight, glycaemic control, blood lipids and blood pressure in type 2 diabetes: meta-analysis of randomised controlled trials. *Br J Nutr* 2013; 110:781–789.
 23. Santesso N, Akl EA, Bianchi M, *et al.* Effects of higher- versus lower-protein diets on health outcomes: a systematic review and meta-analysis. *Eur J Clin Nutr* 2012; 66:780–788.
 24. Cederholm T, Cruz-Jentoft AJ, Maggi S. Sarcopenia and fragility fractures. *Eur J Phys Rehabil Med* 2013; 49:111–117.

25. Paddon-Jones D, Sheffield-Moore M, Urban RJ, *et al.* Essential amino acid and carbohydrate supplementation ameliorates muscle protein loss in humans during 28 days bedrest. *J Clin Endocrinol Metab* 2004; 89:4351–4358.
 26. Ferrando AA, Paddon-Jones D, Hays NP, *et al.* EAA supplementation to increase nitrogen intake improves muscle function during bed rest in the elderly. *Clin Nutr* 2010; 29:18–23.
 27. English KL. Effects of leucine on skeletal muscle during 14 d bed rest in middle-aged adults. Galveston, TX: The University of Texas Medical Branch; 2013.
 28. Kortebein P, Symons TB, Ferrando A, *et al.* Functional impact of 10 days of bed rest in healthy older adults. *J Gerontol A Biol Sci Med Sci* 2008; 63:1076–1081.
 29. Symons TB, Sheffield-Moore M, Mamerow MM, *et al.* The anabolic response to resistance exercise and a protein-rich meal is not diminished by age. *J Nutr Health Aging* 2011; 15:376–381.
 30. English KL, Paddon-Jones D. Protecting muscle mass and function in older adults during bed rest. *Curr Opin Clin Nutr Metab Care* 2010; 13:34–39.
 31. Deutz NE, Pereira SL, Hays NP, *et al.* Effect of β -hydroxy- β -methylbutyrate (HMB) on lean body mass during 10 days of bed rest in older adults. *Clin Nutr* 2013; 32:704–712.
- This study reports that a low volume supplement (HMB) partially preserved muscle mass during physical inactivity. This has potential application for clinical populations and individuals unable to consume sufficient protein during periods of metabolic stress.
32. Casperson SL, Sheffield-Moore M, Hewlings SJ, Paddon-Jones D. Leucine supplementation chronically improves muscle protein synthesis in older adults consuming the RDA for protein. *Clin Nutr* 2012; 31:512–519.
 33. Leidy HJ. Evidence supporting a diet rich in protein to improve appetite control, satiety, and weight management across the lifespan. American Meat Science Association 65th Reciprocal Meat Conference; 17–20 June 2012; North Dakota State University, Fargo, North Dakota. 2012. pp. 1–5.
 34. van der Klaauw A, Keogh J, Henning E, *et al.* High protein intake stimulates GLP1 and PYY release. *Obesity* 2013; 21:1602–1607.
 35. Leidy HJ, Lepping RJ, Savage CR, Harris CT. Neural responses to visual food stimuli after a normal vs. higher protein breakfast in breakfast-skipping teens: a pilot fMRI study. *Obesity (Silver Spring)* 2011; 19:2019–2025.
 36. Leidy HJ, Ortinau LC, Douglas SM, Hoertel HA. Beneficial effects of a higher-protein breakfast on the appetitive, hormonal, and neural signals controlling energy intake regulation in overweight/obese, 'breakfast-skipping,' late-adolescent girls. *Am J Clin Nutr* 2013; 97:677–688.
 37. Leidy HJ, Mattes RD, Campbell WW. Effects of acute and chronic protein intake on metabolism, appetite, and ghrelin during weight loss. *Obesity (Silver Spring)* 2007; 15:1215–1225.
 38. Leidy HJ, Carnell NS, Mattes RD, Campbell WW. Higher protein intake preserves lean mass and satiety with weight loss in preobese and obese women. *Obesity (Silver Spring)* 2007; 15:421–429.
 39. Symons TB, Sheffield-Moore M, Wolfe RR, Paddon-Jones D. A moderate serving of high-quality protein maximally stimulates skeletal muscle protein synthesis in young and elderly subjects. *J Am Diet Assoc* 2009; 109:1582–1586.
 40. Phillips BE, Hill DS, Atherton PJ. Regulation of muscle protein synthesis in humans. *Curr Opin Clin Nutr Metab Care* 2012; 15:58–63.
 41. Yang Y, Breen L, Burd NA, *et al.* Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr* 2012; 108:1780–1788.
 42. Robinson MJ, Burd NA, Breen L, *et al.* Dose-dependent responses of myofibrillar protein synthesis with beef ingestion are enhanced with resistance exercise in middle-aged men. *Appl Physiol Nutr Metab* 2013; 38:120–125.
 43. Leidy HJ, Armstrong CL, Tang M, *et al.* The influence of higher protein intake and greater eating frequency on appetite control in overweight and obese men. *Obesity (Silver Spring)* 2010; 18:1725–1732.
 44. Leidy HJ, Tang M, Armstrong CL, *et al.* The effects of consuming frequent, higher protein meals on appetite and satiety during weight loss in overweight/obese men. *Obesity (Silver Spring)* 2011; 19:818–824.
 45. Arciero P J, Ormsbee M J, Gentile C L, *et al.* Increased protein intake and meal frequency reduces abdominal fat during energy balance and energy deficit. *Obesity (Silver Spring)* 2013; 21:1357–1366.
 46. Leidy HJ. The benefits of breakfast consumption to combat obesity and diabetes in young people. *Am J Lifestyle Med* 2013; 7:98–104.
 47. Leidy HJ, Bossingham MJ, Mattes RD, Campbell WW. Increased dietary protein consumed at breakfast leads to an initial and sustained feeling of fullness during energy restriction compared to other meal times. *Br J Nutr* 2009; 101:798–803.