



## Review

# A review of lifestyle factors that contribute to important pathways associated with major depression: Diet, sleep and exercise



Adrian L. Lopresti <sup>a,\*</sup>, Sean D. Hood <sup>b</sup>, Peter D. Drummond <sup>a</sup>

<sup>a</sup> School of Psychology, Murdoch University, Perth, Western Australia 6150, Australia

<sup>b</sup> School of Psychiatry & Clinical Neurosciences, University of Western Australia, Perth, Western Australia 6009, Australia

## ARTICLE INFO

## Article history:

Received 25 September 2012

Received in revised form

16 January 2013

Accepted 17 January 2013

Available online 14 February 2013

## Keywords:

Depression

Diet

Exercise

Sleep

Physical activity

## ABSTRACT

Research on major depression has confirmed that it is caused by an array of biopsychosocial and lifestyle factors. Diet, exercise and sleep are three such influences that play a significant mediating role in the development, progression and treatment of this condition. This review summarises animal- and human-based studies on the relationship between these three lifestyle factors and major depressive disorder, and their influence on dysregulated pathways associated with depression: namely neurotransmitter processes, immuno-inflammatory pathways, hypothalamic–pituitary–adrenal (HPA) axis disturbances, oxidative stress and antioxidant defence systems, neuroprogression, and mitochondrial disturbances. Increased attention in future clinical studies on the influence of diet, sleep and exercise on major depressive disorder and investigations of their effect on physiological processes will help to expand our understanding and treatment of major depressive disorder. Mental health interventions, taking into account the bidirectional relationship between these lifestyle factors and major depression are also likely to enhance the efficacy of interventions associated with this disorder.

© 2013 Elsevier B.V. All rights reserved.

## Contents

1. Introduction	13
2. Methods	13
2.1. Search strategy	13
2.2. Eligibility criteria	13
3. Dysregulated pathways in major depression	13
3.1. Neurotransmitter imbalances	13
3.2. HPA disturbances	14
3.3. Oxidative & nitrosative stress	14
3.4. Neuroprogression	14
3.5. Mitochondrial disturbances	14
3.6. Immuno-inflammation	14
4. Lifestyle factors associated with major depression	15
4.1. The relationship between diet and depression	15
4.1.1. Diet and its effect on inflammation	15
4.1.2. Diet and its effect on neurotransmitters	16
4.1.3. Diet and its effect on oxidative stress	16
4.1.4. Diet and its effect on neuroprogression	16
4.1.5. Diet and its effect on the HPA axis	16
4.1.6. Diet and its effect on mitochondria	17
4.2. The relationship between sleep and depression	17
4.2.1. Sleep and its effect on inflammation	17
4.2.2. Sleep and its effect on neurotransmitters	17
4.2.3. Sleep and its effect on oxidative stress	17

\* Correspondence to: 4/ 165 Summerlakes, Pde Ballajura, Western Australia 6066, Australia. Tel.: +61 892486904; fax: +61 892484274.  
E-mail address: [a.lopresti@murdoch.edu.au](mailto:a.lopresti@murdoch.edu.au) (A.L. Lopresti).

4.2.4. Sleep and its effect on neuroprogression ..... 18  
 4.2.5. Sleep and its effect on the HPA axis ..... 18  
 4.2.6. Sleep and its effect on mitochondria ..... 18  
 4.3. The relationship between exercise & depression ..... 18  
 4.3.1. Exercise and its effect on inflammation ..... 19  
 4.3.2. Exercise and its effect on neurotransmitters ..... 19  
 4.3.3. Exercise and its effect on oxidative stress ..... 19  
 4.3.4. Exercise and its effect on the HPA axis ..... 19  
 4.3.5. Exercise and its effect on neuroprogression ..... 20  
 4.3.6. Exercise and its effect on mitochondria ..... 20  
 5. Conclusion and directions for future research ..... 20  
 Conflict of interest ..... 21  
 Role of funding source ..... 21  
 Acknowledgements ..... 21  
 References ..... 21

**1. Introduction**

Technological advances have changed according to how we communicate, the activities we engage in, our occupational and recreational pursuits, and even the foods that we eat. While sport and leisure activity levels have remained stable or increased slightly over time, physical activity associated with work, home, and transportation has declined significantly (Brownson et al., 2005; Juneau and Potvin, 2010). In the United States it was estimated that over the past 50 years occupation-related energy expenditure decreased by more than 100 cal/day (Church et al., 2011). Driving to work increased from 67% of the working north American population in 1960 to 88% in 2000 (Brownson et al., 2005) and, in US school children, walking or riding bikes to school decreased from 40% in 1969 to 13% in 2001 (McDonald, 2007). Dietary changes are also significant as worldwide sugar consumption has increased by 74-kcal/day per person from 1962 to 2000. Of this increase, 80% was derived from sugared beverages with additional contributions from restaurant and fast food sources (Popkin and Nielsen, 2003). Alarmingly, sugar consumption has increased most in children aged 6–11 years with an approximate 20% increase from 1988 to 2004 (Wang et al., 2008).

These and other changes of modernity over the past few decades have coincided with a reported increase in the prevalence of many psychiatric problems, including major depression. Between 1991–1992 and 2001–2002, one-year prevalence rates of major depression increased from 3.33% to 7.06% in a community population of American adults (Compton et al., 2006). Increases have also been observed in Australian communities with prevalence rates rising from 6.8% to 10.3% between 1998 and 2008 (Goldney et al., 2010). While these increased rates of depression may be due, in part, to improvements in diagnostic recognition, changes in diagnostic criteria and increased community acceptance of this condition, contemporary lifestyles might also explain why depression is on the rise. However, underlying mechanisms are not well understood. This review provides a summary of three major lifestyle mediators—diet, exercise and sleep-associated with major depression and their impact on a range of relevant biological and physiological pathways.

**2. Methods**

*2.1. Search strategy*

The PubMed, Google Scholar, and PsycInfo databases were searched from all years of record until August 2012. Most references were obtained from combinations of the following key terms:

“depression”, “diet”, “nutrients”, “sleep”, “exercise”, “inflammation”, “oxidative stress”, “mitochondria”, “neurogenesis”, “BDNF”, “HPA”, “cortisol”, “serotonin” and “monoamines”. The reference lists of relevant papers were also examined to locate additional studies that were not identified by the database searches.

*2.2. Eligibility criteria*


Studies were included in this review if they were published in English, comprised animal or human investigations and examined areas of exercise, sleep or diet and their impact either on inflammation, hypothalamic–pituitary–adrenal (HPA) axis, neurotransmitters, neuroprogression and oxidative/nitrosative stress.

**3. Dysregulated pathways in major depression**

Major depression has a multifactorial aetiology arising from environmental, psychological, genetic and biological factors. As outlined in Fig. 1, research over the past decade has clarified that depression is associated with neurotransmitter imbalances, HPA disturbances, dysregulated inflammatory pathways, increased oxidative and nitrosative damage, neuroprogression, and mitochondrial disturbances (Leonard and Maes, 2012; Lopresti et al., 2012; Maes et al., 2009c; Manji et al., 2001; Raison and Miller, 2011). While these disturbances will each be discussed briefly that they are not mutually exclusive.

*3.1. Neurotransmitter imbalances*

Imbalances in the production and transmission of neurotransmitters such as serotonin, dopamine, noradrenaline and glutamate are commonly observed in the central nervous system in major depression (Maletic et al., 2007). Deficiencies in serotonin



**Fig. 1.** Multiple pathways associated with major depression.

availability, the most extensively studied neurotransmitter in depression, is supported by studies using tryptophan depletion models (which reduces central serotonin synthesis) (Hood et al., 2005; Toker et al., 2010) and findings of serotonin receptor abnormalities in depressed patients (Carr and Lucki, 2011). Depression is also associated with an increased availability of monoamine oxidase, an enzyme that metabolises serotonin and other monoamines in the brain (Meyer et al., 2006), and abnormalities in the expression of the enzyme tryptophan hydroxylase, which is involved in serotonin synthesis (Matthes et al., 2010). However, the strongest evidence of neurotransmitter imbalances in depression comes from the popular use and efficacy of pharmaceutical antidepressants, such as selective serotonin reuptake inhibitors (SSRIs), which are thought to alleviate depression by increasing the availability of monoamines such as serotonin, noradrenaline (i.e., nor-epinephrine) and, possibly, dopamine (Connolly and Thase, 2012).

### 3.2. HPA disturbances

Dysfunction in the HPA axis is common in patients with major depression (Pariante and Lightman, 2008). This is characterised by heightened cortisol secretion in patients presenting with melancholic depression, and reduced levels in atypical depression (Gold and Chrousos, 2002). Depression is also associated with hypersecretion of corticotropin-releasing hormone (CRH) and impairment in responsiveness to glucocorticoids (Pariante and Lightman, 2008). An increased size and activity of the pituitary and adrenal glands are also found in major depression (Nemeroff et al., 1992). Successful treatment with antidepressants is associated with a normalisation of HPA axis activity and restoration in glucocorticoid receptor function (Anacker et al., 2011a, 2011b).

### 3.3. Oxidative & nitrosative stress

Decreased antioxidant status and elevated oxidative and nitrosative stress are found in patients with major depression (Maes et al., 2011a). This is evidenced by reduced plasma concentrations of important antioxidants such as vitamin C (Khanzode et al., 2003), vitamin E (Maes et al., 2000; Owen et al., 2005), and coenzyme Q<sub>10</sub> (Maes et al., 2009b), and by reduced antioxidant enzyme activity such as glutathione peroxidase (Maes et al., 2011d). These deficiencies in antioxidant defences impair protection against reactive oxygen species (ROS), leading to damage to fatty acids, proteins and DNA (Maes et al., 2011a).

Depression is also associated with increased levels of lipid peroxidation, comprising elevations in malondialdehyde (Ozcan et al., 2004; Sarandol et al., 2007; Wei et al., 2009), and increased oxidative damage to DNA, characterised by increased levels of 8-hydroxy-2-deoxyguanosine (Forlenza and Miller, 2006; Maes et al., 2009a). Depression is also associated with increased plasma levels of peroxides and xanthine oxidase (Herken et al., 2007; Maes et al., 2010). The efficacy of antioxidant therapies for depression is unknown, although N-acetylcysteine, a powerful antioxidant, was found to be useful for depressive episodes in bipolar disorder (Berk et al., 2008; Magalhaes et al., 2011) and zinc, which serves as a strong antioxidant, also has antidepressant activity (Szewczyk et al., 2011).

### 3.4. Neuroprogression

Neurogenesis and neuronal plasticity are compromised in major depression, with subsequent neurodegeneration (Lee and Kim, 2010). This results in stress-induced alterations to the number and shape of neurons and glia in brain regions of

depressed patients (Duman, 2009) and decreased proliferation of neural stem cells (Eyre and Baune, 2012).

Brain-derived neurotrophic factor (BDNF) is the most abundant and widely distributed neurotrophin in the central nervous system, involved in neuronal survival, growth and proliferation (Martinowich and Lu, 2008). BDNF levels are low in people with major depression (Duman, 2009; Lee and Kim, 2010). However, BDNF levels increase with chronic administration of several classes of antidepressants, including monoamine oxidase inhibitors, SSRIs, tricyclic agents, and SNRIs (Duman and Monteggia, 2006; Sen et al., 2008). Early life and chronic stress, which is often typical in patients with major depression, also has detrimental effects on BDNF (Martinowich et al., 2007; Nagahara and Tuszynski, 2011).

### 3.5. Mitochondrial disturbances


Mitochondria are intracellular organelles that generate most of the cell's supply of adenosine triphosphate (ATP) and are also involved in a range of other processes such as signalling, cellular differentiation, cell death, and the control of the cell cycle and cell growth (McBride et al., 2006). High concentrations of mitochondria are found in the brain which increases its vulnerability to reductions in aerobic metabolism (Pieczenik and Neustadt, 2007).

Depression is associated with mitochondrial dysfunction or disease with evidence of deletions of mitochondrial DNA (Gardner and Boles, 2008a; Shao et al., 2008), and lower activities of respiratory chain enzymes and ATP production (Gardner et al., 2003). Depressed patients presenting with somatic complaints also have low ATP production rates in biopsied muscles (Gardner and Boles, 2008a, 2008b). In addition, rates of depression are increased in patients with mitochondrial disorders (Fattal et al., 2007; Koene et al., 2009).

### 3.6. Immuno-inflammation

Increased inflammation in major depression has been confirmed in three recent meta-analyses. Elevated levels of C-reactive protein (CRP), interleukin-1 (IL-1), and interleukin-6 (IL-6) were reported in a meta-analysis on depression in clinic and community samples (Howren et al., 2009), levels of tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ) and IL-6 were significantly higher in depressed patients than controls (Dowlati et al., 2010), and blood levels of soluble interleukin-2 receptors, TNF- $\alpha$  and IL-6 were higher in a meta-analysis on patients with major depressive disorder than controls (Liu et al., 2012b). Major depression is also characterised by a Th-1-like cell-mediated response, with evidence of increased production of interferon- $\gamma$  (IFN- $\gamma$ ), increased IFN- $\gamma$ /IL-4 ratios and increased neopterin levels (Maes et al., 1994; Myint et al., 2005). In addition, anti-depressant medications have anti-inflammatory effects (Hannestad and Bloch, 2011).

An elevated immuno-inflammatory response in major depression is further supported by investigations into kynurenine pathway metabolites or TRYCATS (tryptophan catabolites along the IDO pathway) (Dantzer et al., 2011; Maes et al., 2011c). As shown in Fig. 2, TRYCATS are produced by the breakdown of tryptophan, involving the enzyme indoleamine 2,3-dioxygenase (IDO). IDO is expressed in multiple cell types including macrophages, dendritic cells, astrocytes and microglia and is strongly activated by the pro-inflammatory cytokine IFN- $\gamma$  and to a lesser extent TNF- $\alpha$ , IL-1, and IL-6. These TRYCATS have both neurotoxic and neuro-protective qualities. Preliminary research has demonstrated a relationship between depression and low levels of the neuroprotective TRYCAT, kynurenine acid (KYNA) (Maes et al., 2011b; Myint et al., 2007; Wichers et al., 2005), and high levels of the excitotoxic TRYCAT, quinolinic acid (QUIN) (Raison et al., 2010a;



**Fig. 2. Kynurenine pathway and its metabolites.** The kynurenine pathway starts with the degradation of tryptophan by the enzyme, indoleamine 2,3-dioxygenase (IDO) which is upregulated by pro-inflammatory cytokines (e.g. IFN- $\gamma$ , TNF- $\alpha$ , IL-, IL-6). These TRYCATS (tryptophan catabolites along the IDO pathway) have neuroprotective and neurotoxic effects on the CNS and influence monoaminergic transmission.

Steiner et al., 2011). However, further research is warranted as Hughes et al. (2012) found no differences in IDO expression or plasma levels of TRYCATS between depressed patients and controls.

#### 4. Lifestyle factors associated with major depression

Dysregulation in the pathways reviewed above can be influenced by environmental, social, psychological, lifestyle, genetic and physiological factors (Hidaka, 2012; Leonard and Maes, 2012). Diet, sleep and exercise are three such influences that play an important role in the aetiology, progression and treatment of depression. A bidirectional relationship likely exists between depression and these mediators.

##### 4.1. The relationship between diet and depression

An association between diet and depression has now been confirmed in prospective and epidemiological studies. For example, in elderly men and women, the consumption of fish, vegetables, olive oil, and cereal correlated negatively with the severity of depressive symptoms (Mamplekou et al., 2010). The benefits from fish and olive oil intake remained significant even when adjusted for confounders such as age, sex, education status, BMI and physical activity status, as well as the presence of a number of medical conditions. In a prospective study, and after adjusting for sex, age, smoking status, BMI, physical activity levels and employment status, adherence to a Mediterranean diet comprising high levels of vegetables, fruit, nuts, cereal, legumes, and fish, a moderate alcohol intake, and a low consumption of meat or meat products and whole-fat dairy, was protective against the development of depression (Sanchez-Villegas et al., 2009). In a study by Jacka et al. (2010b), consuming a 'traditional' diet comprising vegetables, fruit, meat, fish, and whole grains was also associated with a 35% reduced risk of depression or dysthymia. Research into the diet of adolescents (Jacka et al., 2010a) and of low socio-economic, community dwelling, elderly people (German et al., 2011) has also provided evidence for an association between diet quality and depression. Depressive symptoms are also positively associated with the consumption of sweets (Jeffery et al., 2009). Similarly, high intake of fast food (hamburgers, sausages, pizza) and processed pastries (muffins, doughnuts, croissants) are associated with an increased risk of depression up to 6 years later (Sanchez-Villegas et al., 2012).

High-quality treatment studies investigating the impact of diet on depression are scarce, although in a randomised-controlled trial, meat-eating adults placed on a two-week vegetarian diet reported significantly greater improvements in mood compared to participants who continued to eat meat, fish or poultry (Beezhold and Johnston, 2012). In another randomised-controlled trial, six days on a low protein diet significantly decreased depressive symptoms in type 2 diabetics (Ciarambino et al., 2011), and in a randomised study on overweight and obese individuals, those placed on an energy-restricted, low-fat diet for one-year experienced greater improvements in mood compared to participants on an energy-restricted, low-carbohydrate diet (Brinkworth et al., 2009). These changes were independent of weight loss.

Polyunsaturated fatty acids (PUFAs) and particularly omega-3 essential fatty acids ( $\omega$ -3 EFA) have received significant attention in relation to depression. In a meta-analysis of 14 studies comparing the levels of PUFAs between depressed patients and control subjects, the levels of eicosapentaenoic acid (EPA), docosahexaenoic acid (DHA), and total  $\omega$ -3 EFA were significantly lower in depressed patients than controls (corresponding to effect sizes of  $-0.18$ ,  $-0.35$  and  $-0.51$ , respectively). There was no significant change in arachidonic acid (AA) or total  $\omega$ -6 PUFAs (Lin et al., 2010). A meta-analysis of the effects of EPA supplementation in 15 clinical trials in depressed populations revealed beneficial effects from fish oil containing high levels of EPA (effect size=0.53) (Sublette et al., 2011).

Other investigations on the relationship between nutrients and depression have demonstrated a role of folate (Farah, 2009; Gilbody et al., 2007; Morris et al., 2008), tryptophan (Cowen et al., 1989; DeMyer et al., 1981; Maes et al., 1987), zinc (Cope and Levenson, 2010; Lai et al., 2012; Szewczyk et al., 2011), iron (Maes et al., 1996; Stewart and Hirani, 2012; Vahdat Shariatpanaahi et al., 2007; Yi et al., 2011), CoQ10 (Maes et al., 2009b), vitamin B6 (Merete et al., 2008; Moorthy et al., 2012; Skarupski et al., 2010; Williams et al., 2005), vitamin B12 (Hintikka et al., 2003; Moorthy et al., 2012), and selenium (Gao et al., 2012; Mokhber et al., 2011; Pasco et al., 2012). However, findings on most of these nutrients require further investigation before definitive conclusions about their relationship with depression can be made.

##### 4.1.1. Diet and its effect on inflammation

There is now strong evidence in human studies that adherence to a Mediterranean diet is associated with reduced inflammatory markers (Camargo et al., 2012; Richard et al., 2012; Urpi-Sarda et al., 2012). In a study on people with metabolic syndrome, five weeks on a Mediterranean diet corresponded with lowered plasma CRP and an arbitrary inflammatory score that included CRP, IL-6, IL-18, and TNF- $\alpha$ . These changes were independent of any weight loss (Richard et al., 2012). Compared to participants placed on a low-fat diet, one year on a Mediterranean diet was associated with lowered plasma concentrations of IL-6, and two TNF receptors (Urpi-Sarda et al., 2012). In contrast, intercellular adhesion molecule-1 and TNF receptor concentrations were increased in people consuming a low-fat diet. In another intervention study, postprandial inflammatory gene expression in mononuclear cells decreased after three weeks on either a Mediterranean diet enriched with olive oil, a diet rich in saturated fatty acids, or a low-fat/high-carbohydrate diet enriched with  $\omega$ -3 PUFU compared to the other diets (Camargo et al., 2012; Yubero-Serrano et al., 2012). Luciano et al. (2012) also found that CRP levels were lower in an elderly population on a Mediterranean diet compared to a standard 'healthy diet' comprising a high intake of fruits and low consumption of eggs, spirits or liqueurs, and meats such as bacon, pork, lamb, and sausages.

In relation to the anti-inflammatory effect of PUFAs, in a recent review of 26 randomised clinical trials, dietary  $\omega$ -3 EFAs were found to be associated with lower plasma biomarker levels, reflecting lower levels of inflammation and endothelial activation (e.g., IL-6, CRP, TNF- $\alpha$ , sICAM-1 and GM-CSF) in cardiovascular disease and other chronic and acute diseases. Calder (2012) recently concluded that fatty acids were able to partly inhibit a number of aspects of inflammation including leucocyte chemotaxis, adhesion molecule expression and leucocyte-endothelial adhesive interactions, production of eicosanoids from arachidonic acid, production of inflammatory cytokines, and T cell reactivity.

#### 4.1.2. Diet and its effect on neurotransmitters

Diet quality is important in the production of monoamines such as serotonin and dopamine and can influence receptor sensitivity and neurotransmitter transporters. In animal studies, semi-starvation on a high carbohydrate or protein diet affected serotonin turnover in the brain (Schweiger et al., 1989). In addition, the consumption of sugar as part of a meal (Inam et al., 2006) or eating a high-carbohydrate diet (Buwalda et al., 2001) influenced 5-HT<sub>1A</sub> receptor sensitivity; and one week on a high fat, low carbohydrate diet decreased serotonin release in the hypothalamus (Bananas et al., 2009). In contrast, the acute intake of a carbohydrate-rich food increased brain tryptophan and consequent brain serotonin levels (Fernstrom and Wurtman, 1971). This was likely due to carbohydrates acutely increasing brain tryptophan availability compared to other large neutral amino acids (Wurtman and Wurtman, 1995).

Dopaminergic systems are also influenced by diet as the consumption of combinations of dietary fat and sugar reduced D<sub>2</sub> receptor signalling (Pritchett and Hajnal, 2011), the intake of a high fat diet altered dopamine-related gene expression (Lee et al., 2010; Vucetic et al., 2012), a high-fat diet during early life altered biochemical markers of dopamine signalling in the nucleus accumbens (Teegarden et al., 2009), and the long-term consumption of a low protein-high carbohydrate diet decreased D<sub>2</sub> dopamine receptor density (Hamdi et al., 1992). Striatal dopamine levels were also increased in rats supplemented with strawberry, spinach, or vitamin E (Martin et al., 2000).

Other nutrients which are altered in patients with major depression and that can influence neurotransmitter production include tryptophan (and other large neutral amino acids: valine, leucine, isoleucine, phenylalanine and tyrosine) (Maes et al., 2011c; Markus, 2008; Toker et al., 2010), folic acid (Miller, 2008; Stahl, 2008), zinc (Cichy et al., 2009; Szewczyk et al., 2009, 2011), vitamin B12 (Bottiglieri, 1996; Deana et al., 1977), vitamin B6 (Calderon-Guzman et al., 2004; Demisch and Kaczmarczyk, 1991; Hartvig et al., 1995) and iron (Baumgartner et al., 2012; Burhans et al., 2005; Coe et al., 2009). Omega-3 EFAs are also able to modify monoaminergic neurotransmission (Chalon, 2006; Su, 2009).

#### 4.1.3. Diet and its effect on oxidative stress

Given the crucial role that diet plays in antioxidant intake, it comes as no surprise that diet quality influences levels of oxidative stress. In animal studies, rats fed a high-sugar/high-fat diet had increased lipid peroxidation in the brain (Ribeiro et al., 2009; Stranahan et al., 2011), elevated plasma malondialdehyde (MDA) concentrations (a marker of lipid peroxidation) (Panchal et al., 2011) and increased mRNA expression levels of genes involved in ROS production in both the liver and adipose tissue (Matsuzawa-Nagata et al., 2008). In obese adults with metabolic syndrome, reducing energy intake by 2000 kJ, mainly via carbohydrate restriction, was associated with decreased oxidative stress and increased levels of antioxidant markers,

alpha-tocopherol and ceruloplasmin (Skalicky et al., 2009). However, placing adults with metabolic syndrome on a 12-week high-fat diet or low-fat, high complex-carbohydrate diet had no effect on markers of oxidative stress and inflammation (Pettersson et al., 2010).

The Mediterranean diet is associated with increased circulating plasma antioxidant levels and decreased oxidative stress (Azzini et al., 2011; Esposito et al., 2011; Yubero-Serrano et al., 2011). The protective properties of this diet may be derived not only from its increased antioxidant concentration but also through its high raw food intake, lower production of cooking-related oxidants and consequent decreased use of nutritional and endogenous antioxidants, and increased fibre intake (Ghiselli et al., 1997). Olive oil, the main source of fat in the Mediterranean diet, is also effective in lowering lipid peroxidation and oxidative stress (Alarcon de la Lastra et al., 2001; Fito et al., 2007)

#### 4.1.4. Diet and its effect on neuroprogression

Diet quality is important for the brain given its capacity both to enhance neurogenic factors and to influence rates of neurodegeneration. In animal studies, brain levels of BDNF decreased in rats maintained on a high carbohydrate diet (Maioli et al., 2012) and high fat diet (Yamada-Goto et al., 2012). Human trials have also demonstrated a relationship between diet and BDNF. Compared to a low-fat diet, adherence to a Mediterranean diet was associated with an improvement in plasma BDNF concentration in individuals with depression (Sanchez-Villegas et al., 2011). In healthy adults, a high-fat meal decreased plasma BDNF by almost 30% (Karczewska-Kupczewska et al., 2011). The importance of diet on neuroprogression is further confirmed by a study on insulin-resistant, overweight and obese subjects where serum BDNF levels increased after three months on a reduced calorie diet (Araya et al., 2008).

Investigations into the potential effects of  $\omega$ -3 PUFAs have revealed that they may also play a role in neuroprogression. In animal studies,  $\omega$ -3 PUFAs supplementation provided protection against reduced plasticity and normalised BDNF after traumatic brain injury (Wu et al., 2004). During pregnancy and lactation, supplementation with  $\omega$ -3 PUFAs protected levels of BDNF and nerve growth factor (NGF) in female rats when they consumed a micronutrient-imbalanced diet (Sable et al., 2012), while brain levels of BDNF decreased during diets deficient in  $\omega$ -3 PUFAs (Bhatia et al., 2011; Rao et al., 2007). In an human open-label trial, 3 months of  $\omega$ -3 PUFAs supplementation increased serum BDNF levels and prevented posttraumatic distress after accidental injury in patients presenting at an intensive care unit (Matsuoka et al., 2011). However, in a randomised, double-blind, placebo-controlled study of diabetic patients with major depression, 12 weeks of ethyl-EPA supplementation or placebo, in addition to ongoing antidepressant therapy, failed to increase serum BDNF levels (Bot et al., 2011).

#### 4.1.5. Diet and its effect on the HPA axis

Diet composition and timing have a significant influence on acute cortisol secretion due to the primary role of cortisol in gluconeogenesis. However, the long-term effect of diet on HPA activity is not well understood. In a study on women living in a Mediterranean area, a disturbed HPA axis was associated with a higher content of fat and saturated fatty acids in the diet. In contrast, adherence to a dietary pattern closer to the Mediterranean diet was linked with smaller HPA axis disturbances (Garcia-Prieto et al., 2007). Investigations into the effect of  $\omega$ -3 PUFAs on HPA activity have also provided some evidence of their capacity to lower cortisol activity. For example, Delarue et al. (2003) showed that after 3 weeks on a diet supplemented with  $\omega$ -3

PUFAs, levels of plasma noradrenaline and cortisol stimulated by mental stress were significantly blunted. In another study on patients with major depression, serum cortisol levels decreased after 8 weeks of treatment with EPA alone or in combination with fluoxetine (Jazayeri et al., 2010). Finally, intravenous lipopolysaccharide-induced adrenocorticotrophic hormone (ACTH) and cortisol plasma levels decreased significantly in healthy subjects supplemented with one month of fish oil compared to placebo (Michaeli et al., 2007).

#### 4.1.6. Diet and its effect on mitochondria

Mitochondrial dysfunction is influenced significantly by nutrition (Civitaresse et al., 2007; Hepple, 2009; Page et al., 2010; Vitetta and Anton, 2007). Increased fatty acid exposure, resulting from high fat diets or overfeeding, is linked both with decreased mitochondrial number and markers of oxidative phosphorylation. Conversely, caloric restriction can stimulate mitochondrial biogenesis by elevating the transcriptional processes that regulate mitochondrial mass, improve mitochondrial efficiency, activate ROS scavenging mechanisms, and lower ROS production (Civitaresse et al., 2007). Dietary antioxidants or caloric restriction, as well as chemical antioxidants, can lower mitochondrial ROS production (Vitetta and Anton, 2007). Nutrients such as CoQ10, vitamin B2 and L-carnitine also have a significant influence on mitochondrial metabolism (Gardner and Boles, 2011).

## 4.2. The relationship between sleep and depression

In the general population approximately 30% of people report symptoms of insomnia, 10–20% describe dissatisfaction with sleep, and approximately 6% have a formal diagnosis of insomnia (Leger et al., 2000; Ohayon, 2002). These rates are significantly increased in major depression, with as high as 90% of patients reporting sleep disturbances (Motivala et al., 2006; Riemann and Voderholzer, 2003). Insomnia is also one of the most common prodromal features of depression with sleep symptoms preceding an episode of depression in 40% of cases. A history of persistent insomnia is also associated with a significantly increased risk of developing a new depressive episode (Taylor et al., 2005). In a recent meta-analysis, compared to people with no sleep difficulties, non-depressed people with insomnia were predicted to have a twofold increased risk of developing depression (Baglioni et al., 2011). Depressed patients suffering from insomnia also have a poorer response to treatment and are at increased risk of relapse (Dombrovski et al., 2008).

Further support for a relationship between sleep and depression is provided by studies documenting improvements in mood and depressive symptoms following insomnia-specific interventions. For example, in a randomised controlled study, Manber et al. (2008) found that augmenting antidepressant medication with a symptom-focused cognitive-behavioural therapy for insomnia (CBTI) enhanced treatment outcomes in participants with co-morbid major depression and insomnia. Compared to participants allocated to control treatment, people receiving CBTI experienced significantly greater remission rates in both depression (61.5% vs. 33.3%) and insomnia (50% vs. 7.7%). In women with insomnia and breast cancer, CBTI was also more effective in improving sleep, depression and anxiety symptoms than a control condition (Savard et al., 2005). Eight weeks of mindfulness-based cognitive therapy for treating insomnia symptoms also improved sleep, anxiety and depressive symptoms in patients with an anxiety disorder (Yook et al., 2008).

### 4.2.1. Sleep and its effect on inflammation

Sleep difficulties increase inflammatory mediators; conversely elevated inflammatory molecules heighten the risk of sleep problems. In particular, IL-1, IL-6 and TNF- $\alpha$  may be directly involved in sleep regulation (Santos et al., 2007). Data derived from electrophysiological, biochemical and molecular genetic studies demonstrate that these cytokines are sleep regulatory, as they support the regulation of spontaneous sleep-wake behaviour (Opp, 2005). Other cytokines that may also be involved in the regulation of sleep include IL-2, IL-8, IL-15, IL-18, epidermal growth factor, acidic fibroblast growth factor, colony stimulating factor, and interferons (Krueger, 2008).

In patients with major depression, difficulty initiating sleep correlated with increased pre-sleep levels of IL-6 (Motivala et al., 2005). In a study on 210 healthy young men and women, difficulty falling asleep was related to higher morning levels of CRP and IL-6, but only in women (Suarez, 2008). Sleep disturbances also occur in up to 30% of patients with chronic hepatitis C undergoing IFN- $\alpha$  therapy (Sockalingam et al., 2010), and its administration reduces sleep continuity and depth and induces a sleep pattern consistent with insomnia and hyperarousal (Raison et al., 2010b). Studies on patients diagnosed with primary insomnia offer additional evidence for the relationship between sleep and inflammation, as circulating levels of IL-6 and TNF are higher than in healthy sleepers (Burgos et al., 2006; Vgontzas et al., 2002).

Sleep restriction studies in animals and humans provide further confirmation for the association between sleep and inflammation. Although the relationship is not necessarily linear, there is evidence that in humans, sleep restriction increases levels of IL-6, TNF- $\alpha$ , CRP and IL-1 $\beta$  (Motivala, 2011; van Leeuwen et al., 2009; Vgontzas et al., 1999, 2004). Five days of sleep deprivation in healthy adults also modified a number of kynurenine pathway metabolites in healthy adults, including 5-hydroxyindoleacetic acid (5-HIAA), xanthurenic acid and anthranilic acid (Kuhn et al., 1968).

### 4.2.2. Sleep and its effect on neurotransmitters

Surprisingly little research on the influence of sleep on monoamines such as serotonin and dopamine has been conducted, although sleep restriction may disrupt systems associated with monoamine communication. In an animal study, desensitisation in serotonin receptors was detected after eight days of sleep restriction. Despite unlimited recovery sleep, this desensitisation persisted for at least seven days (Roman et al., 2005b), and was independent of adrenal hormones (Roman et al., 2006). In addition, Novati et al. (2008) demonstrated that exposure to a schedule of chronic, partial sleep deprivation reduced sensitivity of 5-HT<sub>1A</sub> receptors and/or receptors for CRH.

### 4.2.3. Sleep and its effect on oxidative stress

It has been proposed that cerebral free radicals accumulate during wakefulness and are removed during sleep (Reimund, 1994). This has been supported by animal studies where sleep loss caused oxidative damage in the brain (Ramanathan et al., 2002; Suer et al., 2011) and increased lipid peroxidation (Thamaraiselvi et al., 2012). However, some animal studies have found that 1–2 weeks of sleep deprivation had no effect on oxidative stress in any brain region, including protein oxidation and lipid peroxidation (D'Almeida et al., 1997; Gopalakrishnan et al., 2004). These inconsistent findings are likely a reflection of differing sleep restriction protocols, and different markers of oxidative stress measured across studies.

Most studies on oxidative stress in clinical sleep research have focused on obstructive sleep apnoea syndrome, which is known

to increase oxidative stress produced by recurrent episodes of ischemia–reperfusion injury (Kent et al., 2011; McNicholas, 2009). However, several studies have also linked oxidative stress with insomnia. For example, levels of thiobarbituric acid reactive substances were elevated in postmenopausal women with insomnia, although blood concentrations of catalase, superoxide dismutase, and glutathione were found to be normal (Hachul de Campos et al., 2006). In an investigation on participants with primary insomnia, significantly lower GSH-Px (selenium-containing antioxidant enzyme) activity and higher MDA levels were found compared with controls (Gulec et al., 2012). Further evidence of a relationship between sleep and oxidative stress is provided by a study revealing increased levels of myeloperoxidase-modified low-density lipoprotein following five nights of sleep restriction in healthy males (Boudjeltia et al., 2011).

#### 4.2.4. Sleep and its effect on neuroprogression

Sleep problems may also contribute to depressive symptomatology via their effect on brain structure, neurogenesis and, in particular, hippocampal function (Lucassen et al., 2010; Meerlo et al., 2009; Novati et al., 2011). Experimental studies show that prolonged sleep restriction or disruption affects hippocampal integrity (Guzman-Marin et al., 2006; Kopp et al., 2006; McDermott et al., 2003; Roman et al., 2005a). For example, in young male rats, one month of chronic sleep restriction reduced dorsal hippocampal volume by 10% (Novati et al., 2011). Hippocampal cell proliferation was also affected by a single day of sleep deprivation in rats (Roman et al., 2005a). Clinical studies have also reported a reduction in hippocampal volume in primary insomnia and sleep apnoea (Morrell et al., 2003; Riemann et al., 2007). Insomnia severity in a sample of patients with post-traumatic stress disorder was also associated with decreased volume in the CA3/dentate hippocampal subfield (Neylan et al., 2010).

While sleep restriction itself may not be neurotoxic, it may enhance neuronal sensitivity to subsequent excitotoxic insults. Novati et al. (2012) found that after 30 days of sleep restriction in rats, there were no adverse effects on cholinergic cells in the nucleus basalis magnocellularis (NBM). However, an injection of a neurotoxic dose of N-methyl-D-aspartate into the NBM caused an accentuated loss of cholinergic NBM cells and cortical fibres in the sleep-restricted rats compared to controls. Thus, chronic sleep restriction may constitute a mild threat to the brain that does not lead to neurodegeneration by itself but increases vulnerability to subsequent neurotoxic challenges.

#### 4.2.5. Sleep and its effect on the HPA axis

Insomnia appears to be associated with hyperarousal. In a study on patients with chronic insomnia, 24-h urinary cortisol levels correlated positively with total wake time. Sleep quality also correlated negatively with urinary levels of catecholamine metabolites, thereby suggesting disturbances in both limbs of the stress system (i.e., the HPA axis and the sympathetic system) (Vgontzas et al., 1998). In another study, ACTH and cortisol secretions were significantly higher in insomniacs compared with normal controls, with greatest elevations in the evening and first half of the night. Cortisol levels were also positively correlated with the severity of reported sleep disturbance (Vgontzas et al., 2001). Increased evening and nocturnal plasma cortisol concentrations were also observed in patients with primary insomnia, with a strong positive correlation between evening cortisol secretion and the number of nocturnal awakenings both in insomniac patients and controls (Rodenbeck and Hajak, 2001).

Findings in sleep deprivation and restriction studies are less consistent, with several studies reporting mild elevations of

cortisol (Leprout et al., 1997; Spiegel et al., 1999) while others have found no change or even slightly decreased levels (Follenius et al., 1992; Kant et al., 1984). In several animal studies, sleep deprivation led to mild activations of the HPA axis and elevated plasma levels of glucocorticoids (Meerlo et al., 2002; Suchecki et al., 1998), whereas others found little or no effect of acute sleep deprivation on glucocorticoid levels (Rechtschaffen et al., 1983). According to Meerlo et al. (2008) the available data from studies in laboratory animals suggest that sleep restriction may gradually change certain brain and neuroendocrine systems in a manner similar to that seen in stress-related disorders such as depression.

#### 4.2.6. Sleep and its effect on mitochondria

Because of the important role of sleep on oxidative stress, it would seem logical to assume that mitochondria will be adversely affected by sleep deprivation. In support of this, sleep deprivation in mice reduced the activity of the complex I, II and III enzymes of the mitochondrial electron transport chain. Complex II and II–III activity was particularly decreased in the hypothalamus of mice during 24-h recovery sleep (Andreazza et al., 2010). In a model developed by Andreazza et al. (2010), it was proposed that sleep restriction may lead to mitochondrial dysfunction which, in turn, increases the production of ROS, leading to increased oxidative damage to lipids, protein and DNA.

#### 4.3. The relationship between exercise & depression

Depression is commonly associated with low levels of physical activity. While data derived from epidemiological and correlational studies do not necessarily confirm causation, a consistent relationship does exist across a number of populations. In adults, an active lifestyle was associated with reduced depressive symptoms independent of education and physical health status. This relationship was stronger in women and those aged 40 years and older (Stephens, 1988). In overweight/obese adults, a reduced risk of depression was associated with increasing moderate-to-vigorous-intensity physical activity and decreasing sedentary time (Vallance et al., 2011). Another study on data from over 4000 men and women aged 20 years or more confirmed that adults with depression spent significantly less time both in light and moderate physical activity than non-depressed adults (Song et al., 2012). In a longitudinal study of over 9000 people, regular physical activity was associated with a reduced likelihood of depressive symptoms at follow-up (Azevedo Da Silva et al., 2012).

Although not as extensive, investigations into sedentary behaviours have also largely confirmed a positive relationship with depression. In a systematic review of seven observational and four intervention studies on adult populations, Teychenne et al. (2010) confirmed that, on balance, sedentary behaviours such as watching television or using the computer were associated with an increased risk of depression. However, evidence was limited by methodological weaknesses in most studies.

The efficacy of exercise as a treatment for depression is summarised in over a dozen recent reviews. In a meta-analysis on supervised and unsupervised physical activity interventions among healthy adults, Conn (2010) concluded that physical activity interventions had a moderate inhibitory effect on depressive symptoms in adults with and without clinical depression (mean effect size of 0.37 for supervised and 0.52 for unsupervised physical activity studies). Carek et al. (2011) maintained from a review of the literature that exercise compared favourably to antidepressant medications as a first-line treatment for mild-to-moderate depression and also improved depressive symptoms when used as an adjunct to medications. Similar antidepressant effects were also found in trials comparing exercise with

cognitive-behavioural therapy (Rimer et al., 2012). Despite these positive findings there is a paucity of research demonstrating long-term beneficial effects of exercise in patients with clinical depression (Krogh et al., 2011).

#### 4.3.1. Exercise and its effect on inflammation

Although a single bout of exercise provokes an acute inflammatory response, primarily in IL-6 (release from muscle increases up to 100-fold during contractile exercise), exercise is followed by an increase in anti-inflammatory cytokines (Pedersen and Fischer, 2007) and a decreased production of the pro-inflammatory cytokines TNF- $\alpha$  and IL-1 $\beta$  (Pedersen et al., 2003). Data also suggest that exercise-induced IL-6 inhibits TNF- $\alpha$  production in the presence of low-grade inflammation (Starkie et al., 2003).

Four recent reviews have primarily revealed anti-inflammatory effects from long-term exercise (Beavers et al., 2010; Mathur and Pedersen, 2008; Ploeger et al., 2009; Thomas and Williams, 2008). In a systematic review of 19 studies on the inflammatory effects of acute and chronic exercise in children and adults, Ploeger et al. (2009) concluded that training programs can attenuate chronic inflammation in some patients with chronic inflammatory disease; however, the exercise training-induced response appeared highly dependent on the type of disease, severity of the disease and the frequency, duration and intensity of the exercise intervention.

Lower inflammatory biomarker concentrations, particularly CRP, and to a lesser extent IL-6, are observed across a wide range of individuals performing more frequent and intense physical activity (Plaisance and Grandjean, 2006; Taaffe et al., 2000). This inverse relationship between CRP and physical activity is consistently seen in men, and to a lesser extent in women (Beavers et al., 2010). In adults, investigations using self-reported measures of physical activity have demonstrated that physically-active individuals have CRP concentrations 19–35% lower than less active individuals (Plaisance and Grandjean, 2006). While these inflammatory markers are attenuated following adjustment for adiposity, a significant relationship between inflammatory biomarkers and physical activity persists (Abramson and Vaccarino, 2002).

In sum, while most studies demonstrate anti-inflammatory effects of exercise in adults and children, the relationship is influenced by a number of factors including the population studied, the type, frequency and duration of exercise, pre-existing medical conditions and initial levels of inflammation. Thus, further studies are required to enable more definitive conclusions, particularly in patients suffering from depression.

#### 4.3.2. Exercise and its effect on neurotransmitters

The antidepressant effects of exercise may be due to its capacity to modify monoamine communication. In animal studies, running increased plasma free tryptophan, brain tryptophan, and levels of the serotonin metabolite, 5-HIAA (Bailey et al., 1993; Chaouloff et al., 1985). Human trials also provide evidence of exercise and its serotonin-enhancing effects. For example, untrained participants randomly assigned to an aerobic exercise group experienced greater changes in serum serotonin levels compared to those in a stretching-control group (Wipfli et al., 2011). Tryptophan availability, the precursor to serotonin, is also increased after acute exercise (Melancon et al., 2012). Three weeks of exercise training also influenced serotonin receptors and serotonin transporters in sedentary males as demonstrated by increased levels of 5-HT transporters (5-HTT) and 5-HT<sub>2A</sub> receptors on isolated platelet membranes. In contrast, four weeks of excessive training in well-trained athletes did not change 5-HTT, and 5-HT<sub>2A</sub> receptor density declined. This suggests that

the impact of exercise on serotonin neurotransmission may depend on the training state of athletes and extent of exertion (Weicker and Struder, 2001).

Exercise is also able to modify dopamine and noradrenergic transmission as evidenced by increased tyrosine hydroxylase expression (Foley and Fleshner, 2008; Kim et al., 2011), elevated striatal dopamine D<sub>2</sub> receptor expression (Vuckovic et al., 2010) and increased noradrenaline levels (Dishman, 1997) in rats exposed to chronic exercise. However, acute exercise comprising 30 min of vigorous exercise in healthy adult volunteers with a history of regular exercise did not change synaptic dopamine concentrations (Wang et al., 2000).

#### 4.3.3. Exercise and its effect on oxidative stress

There is mounting evidence to suggest that exercise is accompanied by an increased generation of free radicals, resulting in measurable elevations in oxidative stress biomarkers after both acute aerobic (Benitez-Sillero et al., 2011; Bloomer, 2008; Fogarty et al., 2011) and anaerobic exercise (Bloomer and Goldfarb, 2004; Pittaluga et al., 2006; Vollaard et al., 2005). Given the substantial evidence for the protective effects of exercise on oxidative stress-associated diseases, this seems paradoxical. However, it is argued that chronic exercise leads to exercise-induced adaptation and resistance (Cooper et al., 2002; Radak et al., 2008). The exercise-induced ROS formation evokes specific adaptation, comprising up-regulation in endogenous antioxidant defences, increased antioxidant/oxidative damage-repairing enzyme activity, increased resistance to oxidative stress, and lowered levels of oxidative damage. This adaptive response seems to be systemic, affecting skeletal muscle, liver, and the brain (Radak et al., 2008). Gomez-Cabrera et al. (2008) argue that because exercise results in an up-regulation of powerful antioxidant enzymes, exercise itself can be considered an antioxidant despite generating free radicals.

#### 4.3.4. Exercise and its effect on the HPA axis

The relationship between exercise and HPA activity is complex as it is influenced by duration, type, intensity and chronicity of exercise; characteristics of the stressor used; and characteristics of the population studied (Campeau et al., 2010; Leal-Cerro et al., 2003; Mastorakos et al., 2005; Stranahan et al., 2008). In an animal study, four weeks of swimming exercise was associated with reduced levels of serum corticosterone and depressive behaviours in rats exposed to high levels of glucocorticoids prenatally (Liu et al., 2012a). In another study, the HPA axis response to lower-intensity stressors decreased in rats exposed to 6 weeks (but not 1 or 3 weeks) of intermittent, voluntary wheel running, although no change occurred following exposure to more intense stressors (Campeau et al., 2010). The complexity of the relationship between exercise and HPA responsiveness was further demonstrated by Droste et al. (2003) who reported that HPA responses in exercising mice were differently influenced by the stressor used and the novelty of the environment.

Investigations into the relationship between exercise and HPA activity in human populations have primarily examined the effect of acute activity on measures such as cortisol and ACTH. Studies on the influence of chronic exercise on HPA activity, and in particular on the HPA response to stressors, are sparse. In general, acute exercise elevates cortisol levels, although this most consistently occurs following moderate-to-severe intensity activity (Hill et al., 2008). In a study on women of varying age and fitness, ACTH levels recovered slowly in older women, particularly those with low fitness levels (Traustadottir et al., 2004). In a study on female adolescents with mild-to-moderate depression, 8 weeks of an exercise regimen improved depressive symptoms and was associated with reductions in 24 h urinary cortisol levels

(Nabkasorn et al., 2006). In another study on patients with chronic low back pain, twelve weeks of high-intensity aerobic exercise was associated with reductions in pain, enhanced mood and improvements in HPA responsiveness to the dexamethasone suppression test (Chatzitheodorou et al., 2008).

#### 4.3.5. Exercise and its effect on neuroprogression

Evidence of the beneficial effects of exercise on brain function is summarised in three recent reviews (Cotman et al., 2007; Dishman et al., 2006; Vivar et al., 2012). Exercise is associated with enhanced adult hippocampal neurogenesis and increased activity-dependent synaptic plasticity. According to Cotman et al. (2007), enhanced hippocampal neurogenesis and increased synaptic plasticity are the most reproducible effects of exercise in the rodent brain. In both young and old animals, exercise stimulated neural progenitor populations, increased the number of new neurons, and promoted survival of these new cells (Brandt et al., 2010; Olson et al., 2006; Wu et al., 2008). In animal studies, exercise also increased BDNF in several brain regions, and there was increased insulin-like growth factor-1 (IGF-1) gene expression and peripheral circulating levels of IGF-1 (Ding et al., 2006; Schwarz et al., 1996). In a review on the effects of exercise on peripheral BDNF in human subjects, it was concluded that exercise temporarily elevated basal BDNF and possibly up-regulated BDNF cellular processing (i.e. synthesis, release, absorption and degradation) (Knaepen et al., 2010). However, there have been no reported findings of long-lasting BDNF responses to acute exercise or training.

Human studies on the effects of exercise-induced BDNF changes in depressed populations are still preliminary, although BDNF levels were transiently increased in elderly women with remitted major depression (Laske et al., 2010) and in unmedicated patients suffering from major depressive disorder (Gustafsson et al., 2009).

#### 4.3.6. Exercise and its effect on mitochondria


An accumulating body of literature has demonstrated that endurance exercise effectively stimulates mitochondrial biogenesis in a wide range of tissues including skeletal muscle, adipose tissue, liver, brain, and kidney (Little et al., 2011). Vina et al. (2009) concluded that exercise, and particularly aerobic exercise, activated mitochondriogenesis in young animals, although its

influence on older animals required further investigation. Eight weeks of treadmill training in a murine model augmented mitochondrial function, as reflected by increased mitochondrial enzyme activities, maximal rate of ATP synthesis in isolated mitochondria, and whole-body maximal O<sub>2</sub> uptake (Chow et al., 2007). Lanza and Nair (2009) reviewed a number of studies which demonstrated that older adults enrolled in exercise training programs respond with increased VO<sub>2</sub> peak, mitochondrial content, oxidative enzyme activities, muscle protein synthesis rates, mitochondrial protein gene transcripts, and mitochondrial DNA copy number.

## 5. Conclusion and directions for future research

While the importance of lifestyle factors such as diet, exercise and sleep are generally acknowledged in the research literature on major depression, the mechanisms of their potential influence are often not fully appreciated. As illustrated in Fig. 3, diet, exercise and sleep can influence several physiological pathways associated with depression. A bi-directional relationship likely exists between depression and these lifestyle factors, thereby creating a potentially increasing cycle of influence. Key symptoms of major depression include changes in appetite, sleep, energy and general motivation levels; all likely to have significant effects on diet, exercise and obviously sleep patterns. The importance of these lifestyle factors was highlighted in a recent paper by Jacka et al. (2012) who argued that depression should be included under the umbrella of non-communicable diseases influenced by lifestyle factors, with increasing efforts directed toward prevention through the promotion of lifestyle changes.

While these lifestyle factors are significant in the aetiology and maintenance of depression, a multitude of other lifestyle influences may also be important. These include chronic stress, social influences, mental and physical effects associated with medical diseases, alcohol and other drug use, chronic pain and even exposure to sunlight/vitamin D. It is these influences, plus a large array of psychological, genetic and biological factors that often make the treatment of depression difficult. Basic interventions comprising attention towards one cause and/or one biochemical mechanism (e.g., targeting a single neurotransmitter disturbance)



**Fig. 3. Potential mechanisms of diet, sleep and exercise on major depression.** Diet, sleep and exercise are associated with depression. These lifestyle factors influence a number of biological processes associated with major depression including neurotransmitter transmission, immuno-inflammation, oxidative and nitrosative stress, HPA balance, neuroprogression and mitochondrial health. Suffering from depression is also likely to lead to changes in diet, sleep and exercise, creating a vicious cycle of change.

makes the goal of remission or recovery less likely. This was highlighted in a recent study where giving simple written recommendations about lifestyle changes for sleep hygiene, physical activity, diet, and sunlight exposure enhanced outcomes to standard antidepressant treatment (Garcia-Toro et al., 2012). Remission/response rates reached 60% in the combined treatment group compared with only 10% in the anti-depressant only group.

Future research needs to be directed toward better understanding the role that diet, exercise, sleep and other lifestyle factors play in depression and other mental health conditions. While this review provides a comprehensive coverage of the research literature, with a specific emphasis on the biological effects of these lifestyle factors on depression, a future more systematic review with well-defined search strategies and inclusion criteria would be beneficial to further elucidate the role these lifestyle factors play in depression, particularly those addressing diet.

A limitation associated with many of the studies reviewed is that significant portions were correlational and/or epidemiological, thereby limiting conclusions about causation. Other influences such as general healthy lifestyle behaviours, socioeconomic status and medical illnesses are examples of confounding factors that may have influenced findings in some studies. Given these and other confounding influences, a significant barrier associated with studies on dietary intervention is being able to accurately provide direct evidence that dietary change causes improvements in mental health. Increasing attention toward randomised, placebo-controlled treatment studies may help to elucidate the mediating roles that these lifestyle factors play in major depression. A paucity of treatment studies has investigated dietary interventions in depressed populations; however, an inherent problem relates to identifying methods to blind such interventions for participants and investigators. While this may be possible with single nutrients or foods, it is likely that the benefits of diet are derived from consuming a range of complementary foods, particularly those characteristic of Mediterranean diets (Ghiselli et al., 1997; Milanese et al., 2011). It is also important that increasing attention be given to measuring changes in important biomarkers associated with inflammation, oxidative stress, HPA regulation, neuroprogression, monoamine and mitochondrial function. Measuring changes in biomarkers and assessing their relationship with affective and behavioural changes should provide a greater understanding of mechanisms of action associated with depression.

#### Conflict of interest

None declared.

#### Role of funding source

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

#### Acknowledgements

Nil

#### References

- Abramson, J.L., Vaccarino, V., 2002. Relationship between physical activity and inflammation among apparently healthy middle-aged and older US adults. *Archives of Internal Medicine* 162, 1286–1292.
- Alarcon de la Lastra, C., Barranco, M.D., Motiva, V., Herreras, J.M., 2001. Mediterranean diet and health: biological importance of olive oil. *Current Pharmaceutical Design* 7, 933–950.
- Anacker, C., Zunszain, P.A., Carvalho, L.A., Pariante, C.M., 2011a. The glucocorticoid receptor: pivot of depression and of antidepressant treatment? *Psychoneuroendocrinology* 36, 415–425.
- Anacker, C., Zunszain, P.A., Cattaneo, A., Carvalho, L.A., Garabedian, M.J., Thuret, S., Price, J., Pariante, C.M., 2011b. Antidepressants increase human hippocampal neurogenesis by activating the glucocorticoid receptor. *Molecular Psychiatry* 16, 738–750.
- Andreazza, A.C., Andersen, M.L., Alvarenga, T.A., De-Oliveira, M.R., Armani, F., Ruiz, F.S., Giglio, L., Moreira, J.C., Kapczinski, F., Tufik, S., 2010. Impairment of the mitochondrial electron transport chain due to sleep deprivation in mice. *Journal of Psychiatric Research* 44, 775–780.
- Araya, A.V., Orellana, X., Espinoza, J., 2008. Evaluation of the effect of caloric restriction on serum BDNF in overweight and obese subjects: preliminary evidences. *Endocrine* 33, 300–304.
- Azevedo Da Silva, M., Singh-Manoux, A., Brunner, E.J., Kaffashian, S., Shipley, M.J., Kivimäki, M., Nabi, H., 2012. Bidirectional association between physical activity and symptoms of anxiety and depression: the Whitehall II study. *European Journal of Epidemiology* 27, 537–546.
- Azzini, E., Polito, A., Fumagalli, A., Intorre, F., Venneria, E., Durazzo, A., Zaccaria, M., Ciarapica, D., Foddai, M.S., Mauro, B., Raguzzini, A., Palomba, L., Maiani, G., 2011. Mediterranean diet effect: an Italian picture. *Nutrition Journal* 10, 125.
- Baglioni, C., Battagliese, G., Feige, B., Spiegelhalter, K., Nissen, C., Voderholzer, U., Lombardo, C., Riemann, D., 2011. Insomnia as a predictor of depression: a meta-analytic evaluation of longitudinal epidemiological studies. *Journal of Affective Disorders* 135, 10–19.
- Bailey, S.P., Davis, J.M., Ahlborn, E.N., 1993. Neuroendocrine and substrate responses to altered brain 5-HT activity during prolonged exercise to fatigue. *Journal of Applied Physiology* 74, 3006–3012.
- Banas, S.M., Rouch, C., Kassis, N., Markaki, E.M., Gerozissis, K., 2009. A dietary fat excess alters metabolic and neuroendocrine responses before the onset of metabolic diseases. *Cellular and Molecular Neurobiology* 29, 157–168.
- Baumgartner, J., Smuts, C.M., Malan, L., Arnold, M., Yee, B.K., Bianco, L.E., Boekschoten, M.V., Muller, M., Langhans, W., Hurrell, R.F., Zimmermann, M.B., 2012. In male rats with concurrent iron and (n-3) fatty acid deficiency, provision of either iron or (n-3) fatty acids alone alters monoamine metabolism and exacerbates the cognitive deficits associated with combined deficiency. *Journal of Nutrition* 142, 1472–1478.
- Beavers, K.M., Brinkley, T.E., Nicklas, B.J., 2010. Effect of exercise training on chronic inflammation. *Clinica Chimica Acta* 411, 785–793.
- Beezhold, B.L., Johnston, C.S., 2012. Restriction of meat, fish, and poultry in omnivores improves mood: a pilot randomized controlled trial. *Nutrition Journal* 11, 9.
- Benitez-Sillero, J.D., Perez-Navero, J.L., Tasset, I., Guillen-Del Castillo, M., Gil-Campos, M., Tunes, I., 2011. Cardiorespiratory fitness and oxidative stress: effect of acute maximal aerobic exercise in children and adolescents. *Journal of Sports Medicine and Physical Fitness* 51, 204–210.
- Berk, M., Copolov, D.L., Dean, O., Lu, K., Jeavons, S., Schapkaitz, I., Anderson-Hunt, M., Bush, A.I., 2008. N-acetyl cysteine for depressive symptoms in bipolar disorder—a double-blind randomized placebo-controlled trial. *Biological Psychiatry* 64, 468–475.
- Bhatia, H.S., Agrawal, R., Sharma, S., Huo, Y.X., Ying, Z., Gomez-Pinilla, F., 2011. Omega-3 fatty acid deficiency during brain maturation reduces neuronal and behavioral plasticity in adulthood. *PLoS One* 6, e28451.
- Bloomer, R.J., 2008. Effect of exercise on oxidative stress biomarkers. *Advances in Clinical Chemistry* 46, 1–50.
- Bloomer, R.J., Goldfarb, A.H., 2004. Anaerobic exercise and oxidative stress: a review. *Canadian Journal of Applied Physiology* 29, 245–263.
- Bot, M., Pouwer, F., Assies, J., Jansen, E.H., Beekman, A.T., De Jonge, P., 2011. Supplementation with eicosapentaenoic omega-3 fatty acid does not influence serum brain-derived neurotrophic factor in diabetes mellitus patients with major depression: a randomized controlled pilot study. *Neuropsychobiology* 63, 219–223.
- Bottiglieri, T., 1996. Folate, vitamin B12, and neuropsychiatric disorders. *Nutrition Reviews* 54, 382–390.
- Boudjeltia, K.Z., Faraut, B., Esposito, M.J., Stenuit, P., Dyzma, M., Antwerpen, P.V., Brohée, D., Vanhamme, L., Moguilevsky, N., Vanhaeverbeek, M., Kerkhofs, M., 2011. Temporal dissociation between myeloperoxidase (MPO)-modified LDL and MPO elevations during chronic sleep restriction and recovery in healthy young men. *PLoS One* 6, e28230.
- Brandt, M.D., Maass, A., Kempermann, G., Storch, A., 2010. Physical exercise increases Notch activity, proliferation and cell cycle exit of type-3 progenitor cells in adult hippocampal neurogenesis. *European Journal of Neuroscience* 32, 1256–1264.
- Brinkworth, G.D., Buckley, J.D., Noakes, M., Clifton, P.M., Wilson, C., 2009. Long-term effects of a very low-carbohydrate diet and a low-fat diet on mood and cognitive function. *Archives of Internal Medicine* 169, 1873–1880.
- Brownson, R.C., Boehmer, T.K., Luke, D.A., 2005. Declining rates of physical activity in the United States: what are the contributors? *Annual Review of Public Health* 26, 421–443.
- Burgos, I., Richter, L., Klein, T., Fiebich, B., Feige, B., Lieb, K., Voderholzer, U., Riemann, D., 2006. Increased nocturnal interleukin-6 excretion in patients with primary insomnia: a pilot study. *Brain, Behavior, and Immunity* 20, 246–253.
- Burhans, M.S., Dailey, C., Beard, Z., Wiesinger, J., Murray-Kolb, L., Jones, B.C., Beard, J.L., 2005. Iron deficiency: differential effects on monoamine transporters. *Nutritional Neuroscience* 8, 31–38.
- Buwalda, B., Blom, W.A., Koolhaas, J.M., van Dijk, G., 2001. Behavioral and physiological responses to stress are affected by high-fat feeding in male rats. *Physiology & Behavior* 73, 371–377.
- Calder, P.C., 2012. Omega-3 polyunsaturated fatty acids and inflammatory processes: nutrition or pharmacology? *British Journal of Clinical Pharmacology*. <http://dx.doi.org/10.1111/j.1365-2125.2012.04374.x>, Epub ahead of print.

- Calderon-Guzman, D., Hernandez-Islas, J.L., Espitia-Vazquez, I., Barragan-Mejia, G., Hernandez-Garcia, E., Santamaria-del Angel, D., Juarez-Olguin, H., 2004. Pyridoxine, regardless of serotonin levels, increases production of 5-hydroxytryptophan in rat brain. *Archives of Medical Research* 35, 271–274.
- Camargo, A., Delgado-Lista, J., Garcia-Rios, A., Cruz-Teno, C., Yubero-Serrano, E.M., Perez-Martinez, P., Gutierrez-Mariscal, F.M., Lora-Aguilar, P., Rodriguez-Cantalejo, F., Fuentes-Jimenez, F., Tinahones, F.J., Malagon, M.M., Perez-Jimenez, F., Lopez-Miranda, J., 2012. Expression of proinflammatory, proatherogenic genes is reduced by the Mediterranean diet in elderly people. *British Journal of Nutrition* 108, 500–508.
- Campeau, S., Nyhuis, T.J., Sasse, S.K., Kryskow, E.M., Herlihy, L., Masini, C.V., Babb, J.A., Greenwood, B.N., Fleshner, M., Day, H.E., 2010. Hypothalamic pituitary adrenal axis responses to low-intensity stressors are reduced after voluntary wheel running in rats. *Journal of Neuroendocrinology* 22, 872–888.
- Carek, P.J., Laibstain, S.E., Carek, S.M., 2011. Exercise for the treatment of depression and anxiety. *International Journal of Psychiatry in Medicine* 41, 15–28.
- Carr, G.V., Lucki, I., 2011. The role of serotonin receptor subtypes in treating depression: a review of animal studies. *Psychopharmacology* 213, 265–287.
- Chalon, S., 2006. Omega-3 fatty acids and monoamine neurotransmission. *Prostaglandins, Leukotrienes and Essential Fatty Acids* 75, 259–269.
- Chaouloff, F., Elghozi, J.L., Guezennec, Y., Laude, D., 1985. Effects of conditioned running on plasma, liver and brain tryptophan and on brain 5-hydroxytryptamine metabolism of the rat. *British Journal of Pharmacology* 86, 33–41.
- Chatzitheodorou, D., Mavromoustakos, S., Milioti, S., 2008. The effect of exercise on adrenocortical responsiveness of patients with chronic low back pain, controlled for psychological strain. *Clinical Rehabilitation* 22, 319–328.
- Chow, L.S., Greenlund, L.J., Asmang, Y.W., Short, K.R., McCrady, S.K., Levine, J.A., Nair, K.S., 2007. Impact of endurance training on murine spontaneous activity, muscle mitochondrial DNA abundance, gene transcripts, and function. *Journal of Applied Physiology* 102, 1078–1089.
- Church, T.S., Thomas, D.M., Tudor-Locke, C., Katzmarzyk, P.T., Earnest, C.P., Rodarte, R.Q., Martin, C.K., Blair, S.N., Bouchard, C., 2011. Trends over 5 decades in US occupation-related physical activity and their associations with obesity. *PLoS One* 6, e19657.
- Ciarambino, T., Ferrara, N., Castellino, P., Paolisso, G., Coppola, L., Giordano, M., 2011. Effects of a 6-days-a-week low protein diet regimen on depressive symptoms in young-old type 2 diabetic patients. *Nutrition* 27, 46–49.
- Cichy, A., Sowa-Kucma, M., Legutko, B., Pomierny-Chamiolo, L., Siwek, A., Piotrowska, A., Szewczyk, B., Poleszak, E., Pilc, A., Nowak, G., 2009. Zinc-induced adaptive changes in NMDA/glutamatergic and serotonergic receptors. *Pharmacological Reports* 61, 1184–1191.
- Civitarese, A.E., Smith, S.R., Ravussin, E., 2007. Diet, energy metabolism and mitochondrial biogenesis. *Current Opinion in Clinical Nutrition and Metabolic Care* 10, 679–687.
- Coe, C.L., Lubach, G.R., Bianco, L., Beard, J.L., 2009. A history of iron deficiency anemia during infancy alters brain monoamine activity later in juvenile monkeys. *Developmental Psychobiology* 51, 301–309.
- Compton, W.M., Conway, K.P., Stinson, F.S., Grant, B.F., 2006. Changes in the prevalence of major depression and comorbid substance use disorders in the United States between 1991–1992 and 2001–2002. *American Journal of Psychiatry* 163, 2141–2147.
- Conn, V.S., 2010. Depressive symptom outcomes of physical activity interventions: meta-analysis findings. *Annals of Behavioral Medicine* 39, 128–138.
- Connolly, K.R., Thase, M.E., 2012. Emerging drugs for major depressive disorder. *Expert Opinion on Emerging Drugs* 17, 105–126.
- Cooper, C.E., Vollaard, N.B., Choueiri, T., Wilson, M.T., 2002. Exercise, free radicals and oxidative stress. *Biochemical Society Transactions* 30, 280–285.
- Cope, E.C., Levenson, C.W., 2010. Role of zinc in the development and treatment of mood disorders. *Current Opinion in Clinical Nutrition and Metabolic Care* 13, 685–689.
- Cotman, C.W., Berchtold, N.C., Christie, L.-A., 2007. Exercise builds brain health: key roles of growth factor cascades and inflammation. *Trends in Neurosciences* 30, 464–472.
- Cowen, P.J., Parry-Billings, M., Newsholme, E.A., 1989. Decreased plasma tryptophan levels in major depression. *Journal of Affective Disorders* 16, 27–31.
- D'Almeida, V., Hipolide, D.C., Azzalis, L.A., Lobo, L.L., Junqueira, V.B., Tufik, S., 1997. Absence of oxidative stress following paradoxical sleep deprivation in rats. *Neuroscience Letters* 235, 25–28.
- Dantzer, R., O'Connor, J.C., Lawson, M.A., Kelley, K.W., 2011. Inflammation-associated depression: from serotonin to kynurenine. *Psychoneuroendocrinology* 36, 426–436.
- Deana, R., Vincenti, E., Deana, A.D., 1977. Levels of neurotransmitters in brain of vitamin B12 deficient rats. *International Journal of Vitamin and Nutrition Research* 47, 119–122.
- Delarue, J., Matzinger, O., Binnert, C., Schneider, P., Chiolerio, R., Tappy, L., 2003. Fish oil prevents the adrenal activation elicited by mental stress in healthy men. *Diabetes & Metabolism* 29, 289–295.
- Demisch, L., Kaczmarczyk, P., 1991. Tryptophan metabolism in healthy subjects: influence of pyridoxine after single or repeated administrations. *Advances in Experimental Medicine and Biology* 294, 519–522.
- DeMyer, M.K., Shea, P.A., Hendrie, H.C., Yoshimura, N.N., 1981. Plasma tryptophan and five other amino acids in depressed and normal subjects. *Archives of General Psychiatry* 38, 642–646.
- Ding, Q., Vaynman, S., Akhavan, M., Ying, Z., Gomez-Pinilla, F., 2006. Insulin-like growth factor I interfaces with brain-derived neurotrophic factor-mediated synaptic plasticity to modulate aspects of exercise-induced cognitive function. *Neuroscience* 140, 823–833.
- Dishman, R.K., 1997. Brain monoamines, exercise, and behavioral stress: animal models. *Medicine and Science in Sports and Exercise* 29, 63–74.
- Dishman, R.K., Berthoud, H.R., Booth, F.W., Cotman, C.W., Edgerton, V.R., Fleshner, M.R., Gandevia, S.C., Gomez-Pinilla, F., Greenwood, B.N., Hillman, C.H., Kramer, A.F., Levin, B.E., Moran, T.H., Russo-Neustadt, A.A., Salamone, J.D., Van Hooissen, J.D., Wade, C.E., York, D.A., Zigmond, M.J., 2006. *Neurobiology of Exercise*. Obesity 14, 345–356.
- Dombrowski, A.Y., Cyranowski, J.M., Mulsant, B.H., Houck, P.R., Buysse, D.J., Andreescu, C., Thase, M.E., Mallinger, A.G., Frank, E., 2008. Which symptoms predict recurrence of depression in women treated with maintenance interpersonal psychotherapy? *Depression and Anxiety* 25, 1060–1066.
- Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E.K., Lanctot, K.L., 2010. A meta-analysis of cytokines in major depression. *Biological Psychiatry* 67, 446–457.
- Droste, S.K., Gesing, A., Ulbricht, S., Muller, M.B., Linthorst, A.C., Reul, J.M., 2003. Effects of long-term voluntary exercise on the mouse hypothalamic-pituitary-adrenocortical axis. *Endocrinology* 144, 3012–3023.
- Duman, R.S., 2009. Neuronal damage and protection in the pathophysiology and treatment of psychiatric illness: stress and depression. *Dialogues in Clinical Neuroscience* 11, 239–255.
- Duman, R.S., Monteggia, L.M., 2006. A neurotrophic model for stress-related mood disorders. *Biological Psychiatry* 59, 1116–1127.
- Esposito, K., Di Palo, C., Maiorino, M.I., Petrizzo, M., Bellastella, G., Siniscalchi, I., Giugliano, D., 2011. Long-term effect of Mediterranean-style diet and calorie restriction on biomarkers of longevity and oxidative stress in overweight men. *Cardiology Research and Practice* 2011, 293916.
- Eyre, H., Baune, B.T., 2012. Neuroplastic changes in depression: a role for the immune system. *Psychoneuroendocrinology* 37, 1397–1416.
- Farah, A., 2009. The role of L-methylfolate in depressive disorders. *CNS Spectrums* 14, 2–7.
- Fattal, O., Link, J., Quinn, K., Cohen, B.H., Franco, K., 2007. Psychiatric comorbidity in 36 adults with mitochondrial cytopathies. *CNS Spectrums* 12, 429–438.
- Fernstrom, J.D., Wurtman, R.J., 1971. Brain serotonin content: increase following ingestion of carbohydrate diet. *Science* 174, 1023–1025.
- Fito, M., de la Torre, R., Covas, M.I., 2007. Olive oil and oxidative stress. *Molecular Nutrition & Food Research* 51, 1215–1224.
- Fogarty, M.C., Hughes, C.M., Burke, G., Brown, J.C., Trinick, T.R., Duly, E., Bailey, D.M., Davison, G.W., 2011. Exercise-induced lipid peroxidation: implications for deoxyribonucleic acid damage and systemic free radical generation. *Environmental and Molecular Mutagenesis* 52, 35–42.
- Foley, T.E., Fleshner, M., 2008. Neuroplasticity of dopamine circuits after exercise: implications for central fatigue. *NeuroMolecular Medicine* 10, 67–80.
- Follenius, M., Brandenberger, G., Badesapt, J.J., Libert, J.P., Ehrhart, J., 1992. Nocturnal cortisol release in relation to sleep structure. *Sleep* 15, 21–27.
- Forlenza, M.J., Miller, G.E., 2006. Increased serum levels of 8-hydroxy-2'-deoxyguanosine in clinical depression. *Psychosomatic Medicine* 68, 1–7.
- Gao, S., Jin, Y., Unverzagt, F.W., Liang, C., Hall, K.S., Cao, J., Ma, F., Murrell, J.R., Cheng, Y., Li, P., Bian, J., Hendrie, H.C., 2012. Selenium level and depressive symptoms in a rural elderly Chinese cohort. *BMC Psychiatry* 12, 72.
- Garcia-Prieto, M.D., Tebar, F.J., Nicolas, F., Larque, E., Zamora, S., Garaulet, M., 2007. Cortisol secretory pattern and glucocorticoid feedback sensitivity in women from a Mediterranean area: relationship with anthropometric characteristics, dietary intake and plasma fatty acid profile. *Clinical Endocrinology* 66, 185–191.
- Garcia-Toro, M., Ibarra, O., Gili, M., Serrano, M.J., Olivan, B., Vicens, E., Roca, M., 2012. Four hygienic-dietary recommendations as add-on treatment in depression: a randomized-controlled trial. *Journal of Affective Disorders* 140, 200–203.
- Gardner, A., Boles, R.G., 2008a. Mitochondrial energy depletion in depression with somatization. *Psychotherapy and Psychosomatics* 77, 127–129.
- Gardner, A., Boles, R.G., 2008b. Symptoms of somatization as a rapid screening tool for mitochondrial dysfunction in depression. *BioPsychoSocial Medicine* 2, 7.
- Gardner, A., Boles, R.G., 2011. Beyond the serotonin hypothesis: mitochondria, inflammation and neurodegeneration in major depression and affective spectrum disorders. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 35, 730–743.
- Gardner, A., Johansson, A., Wibom, R., Nennesmo, I., von Döbeln, U., Hagenfeldt, L., Hallstrom, T., 2003. Alterations of mitochondrial function and correlations with personality traits in selected major depressive disorder patients. *Journal of Affective Disorders* 76, 55–68.
- German, L., Kahana, C., Rosenfeld, V., Zabrowsky, I., Wiezer, Z., Fraser, D., Shahar, D.R., 2011. Depressive symptoms are associated with food insufficiency and nutritional deficiencies in poor community-dwelling elderly people. *Journal of Nutrition Health and Aging* 15, 3–8.
- Ghiselli, A., D'Amicis, A., Giacosa, A., 1997. The antioxidant potential of the Mediterranean diet. *European Journal of Cancer Prevention* 6 (Suppl. 1), S15–S19.
- Gilbody, S., Lightfoot, T., Sheldon, T., 2007. Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity. *Journal of Epidemiology and Community Health* 61, 631–637.
- Gold, P.W., Chrousos, G.P., 2002. Organization of the stress system and its dysregulation in melancholic and atypical depression: high vs. low CRH/NE states. *Molecular Psychiatry* 7, 254–275.

- Goldney, R.D., Eckert, K.A., Hawthorne, G., Taylor, A.W., 2010. Changes in the prevalence of major depression in an Australian community sample between 1998 and 2008. *Australian and New Zealand Journal of Psychiatry* 44, 901–910.
- Gomez-Cabrera, M.C., Domenech, E., Viña, J., 2008. Moderate exercise is an antioxidant: upregulation of antioxidant genes by training. *Free Radical Biology and Medicine* 44, 126–131.
- Gopalakrishnan, A., Ji, L.L., Cirelli, C., 2004. Sleep deprivation and cellular responses to oxidative stress. *Sleep* 27, 27–35.
- Gulec, M., Ozkol, H., Selvi, Y., Tuluze, Y., Aydin, A., Besiroglu, L., Ozdemir, P.G., 2012. Oxidative stress in patients with primary insomnia. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 37, 247–251.
- Gustafsson, G., Lira, C.M., Johansson, J., Wisén, A., Wohlfart, B., Ekman, R., Westrin, A., 2009. The acute response of plasma brain-derived neurotrophic factor as a result of exercise in major depressive disorder. *Psychiatry Research* 169, 244–248.
- Guzman-Marin, R., Ying, Z., Sunstova, N., Methippara, M., Bashir, T., Szymusiak, R., Gomez-Pinilla, F., McGinty, D., 2006. Suppression of hippocampal plasticity-related gene expression by sleep deprivation in rats. *Journal of Physiology* 575, 807–819.
- Hachul de Campos, H., Brandao, L.C., D'Almeida, V., Grego, B.H., Bittencourt, L.R., Tufik, S., Baracat, E.C., 2006. Sleep disturbances, oxidative stress and cardiovascular risk parameters in postmenopausal women complaining of insomnia. *Climacteric* 9, 312–319.
- Hamdi, A., Onaivi, E.S., Prasad, C., 1992. A low protein–high carbohydrate diet decreases D2 dopamine receptor density in rat brain. *Life Sciences* 50, 1529–1534.
- Hannestad, J.N.D., Bloch, M., 2011. The effect of antidepressant medication treatment on serum levels of inflammatory cytokines: a meta-analysis. *Neuropsychopharmacology* 36, 2452–2459.
- Hartvig, P., Lindner, K.J., Bjurling, P., Laengstrom, B., Tedroff, J., 1995. Pyridoxine effect on synthesis rate of serotonin in the monkey brain measured with positron emission tomography. *Journal of Neural Transmission. General Section* 102, 91–97.
- Hepple, R.T., 2009. Why eating less keeps mitochondria working in aged skeletal muscle. *Exercise and Sport Sciences Reviews* 37, 23–28.
- Herken, H., Gurel, A., Selek, S., Armutcu, F., Ozen, M.E., Bulut, M., Kap, O., Yumru, M., Savas, H.A., Akyol, O., 2007. Adenosine deaminase, nitric oxide, superoxide dismutase, and xanthine oxidase in patients with major depression: impact of antidepressant treatment. *Archives of Medical Research* 38, 247–252.
- Hidaka, B.H., 2012. Depression as a disease of modernity: explanations for increasing prevalence. *Journal of Affective Disorders* 140, 205–214.
- Hill, E.E., Zack, E., Battaglini, C., Viru, M., Viru, A., Hackney, A.C., 2008. Exercise and circulating cortisol levels: the intensity threshold effect. *Journal of Endocrinological Investigation* 31, 587–591.
- Hintikka, J., Tolmunen, T., Tanskanen, A., Viinamäki, H., 2003. High vitamin B12 level and good treatment outcome may be associated in major depressive disorder. *BMC Psychiatry* 3, 17.
- Hood, S.D., Bell, C.J., Nutt, D.J., 2005. Acute tryptophan depletion. Part I: rationale and methodology. *Australian and New Zealand Journal of Psychiatry* 39, 558–564.
- Howren, M.B., Lamkin, D.M., Suls, J., 2009. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosomatic Medicine* 71, 171–186.
- Hughes, M.M., Carballo, A., McLoughlin, D.M., Amico, F., Harkin, A., Frodl, T., Connor, T.J., 2012. Tryptophan depletion in depressed patients occurs independent of kynurenine pathway activation. *Brain, Behavior, and Immunity* 26, 979–987.
- Inam, Q.U., Haleem, M.A., Haleem, D.J., 2006. Effects of long term consumption of sugar as part of meal on serotonin-1a receptor dependent responses. *Pakistan Journal of Pharmaceutical Sciences* 19, 94–98.
- Jacka, F.N., Kremer, P.J., Leslie, E.R., Berk, M., Patton, G.C., Toumbourou, J.W., Williams, J.W., 2010a. Associations between diet quality and depressed mood in adolescents: results from the Australian Healthy Neighbourhoods Study. *Australian and New Zealand Journal of Psychiatry* 44, 435–442.
- Jacka, F.N., Pasco, J.A., Mykletun, A., Williams, L.J., Hodge, A.M., O'Reilly, S.L., Nicholson, G.C., Kotowicz, M.A., Berk, M., 2010b. Association of Western and traditional diets with depression and anxiety in women. *American Journal of Psychiatry* 167, 305–311.
- Jacka, F.N., Mykletun, A., Berk, M., 2012. Moving towards a population health approach to the primary prevention of common mental disorders. *BMC Medicine* 10, 149.
- Jazayeri, S., Keshavarz, S.A., Tehrani-Doost, M., Djalali, M., Hosseini, M., Amini, H., Chamari, M., Djazayeri, A., 2010. Effects of eicosapentaenoic acid and fluoxetine on plasma cortisol, serum interleukin-1beta and interleukin-6 concentrations in patients with major depressive disorder. *Psychiatric Research* 178, 112–115.
- Jeffery, R.W., Linde, J.A., Simon, G.E., Ludman, E.J., Rohde, P., Ichikawa, L.E., Finch, E.A., 2009. Reported food choices in older women in relation to body mass index and depressive symptoms. *Appetite* 52, 238–240.
- Juneau, C.E., Potvin, L., 2010. Trends in leisure-, transport-, and work-related physical activity in Canada 1994–2005. *Preventive Medicine* 51, 384–386.
- Kant, G.J., Genser, S.G., Thorne, D.R., Pfalser, J.L., Mougey, E.H., 1984. Effects of 72 h sleep deprivation on urinary cortisol and indices of metabolism. *Sleep* 7, 142–146.
- Karczewska-Kupczewska, M., Kowalska, I., Nikołaćuk, A., Adamska, A., Zielińska, M., Kamińska, N., Otziomek, E., Górski, M., Strączkowski, M., 2011. Circulating brain-derived neurotrophic factor concentration is downregulated by intralipid/heparin infusion or high-fat meal in young healthy male subjects. *Diabetes Care* 35, 358–362.
- Kent, B.D., Ryan, S., McNicholas, W.T., 2011. Obstructive sleep apnea and inflammation: relationship to cardiovascular co-morbidity. *Respiratory Physiology & Neurobiology* 178, 475–481.
- Khanzode, S.D., Dakhale, G.N., Khanzode, S.S., Saoji, A., Palasodkar, R., 2003. Oxidative damage and major depression: the potential antioxidant action of selective serotonin re-uptake inhibitors. *Redox Report* 8, 365–370.
- Kim, H., Heo, H.I., Kim, D.H., Ko, I.G., Lee, S.S., Kim, S.E., Kim, B.K., Kim, T.W., Ji, E.S., Kim, J.D., Shin, M.S., Choi, Y.W., Kim, C.J., 2011. Treadmill exercise and methylphenidate ameliorate symptoms of attention deficit/hyperactivity disorder through enhancing dopamine synthesis and brain-derived neurotrophic factor expression in spontaneous hypertensive rats. *Neuroscience Letters* 504, 35–39.
- Knaepen, K., Goekint, M., Heyman, E.M., Meeusen, R., 2010. Neuroplasticity-exercise-induced response of peripheral brain-derived neurotrophic factor: a systematic review of experimental studies in human subjects. *Sports Medicine* 40, 765–801.
- Koene, S., Kozicz, T.L., Rodenburg, R.J., Verhaak, C.M., De Vries, M.C., Wortmann, S., van de Heuvel, L., Smeitink, J.A., Morava, E., 2009. Major depression in adolescent children consecutively diagnosed with mitochondrial disorder. *Journal of Affective Disorders* 114, 327–332.
- Kopp, C., Longordo, F., Nicholson, J.R., Luthi, A., 2006. Insufficient sleep reversibly alters bidirectional synaptic plasticity and NMDA receptor function. *Journal of Neuroscience* 26, 12456–12465.
- Krogh, J., Nordentoft, M., Sterne, J.A., Lawlor, D.A., 2011. The effect of exercise in clinically depressed adults: systematic review and meta-analysis of randomized controlled trials. *Journal of Clinical Psychiatry* 72, 529–538.
- Krueger, J.M., 2008. The role of cytokines in sleep regulation. *Current Pharmaceutical Science* 14, 3408–3416.
- Kuhn, E., Rysanek, K., Brodan, V., 1968. Alterations of tryptophan metabolism induced by sleep deprivation. *Experientia* 24, 901–902.
- Lai, J., Moxey, A., Nowak, G., Vashum, K., Bailey, K., McEvoy, M., 2012. The efficacy of zinc supplementation in depression: systematic review of randomised controlled trials. *Journal of Affective Disorders* 136, e31–e39.
- Lanza, I.R., Nair, K.S., 2009. Muscle mitochondrial changes with aging and exercise. *American Journal of Clinical Nutrition* 89, 467S–471S.
- Laske, C., Banschbach, S., Stransky, E., Bosch, S., Straten, G., Machann, J., Fritsche, A., Hipp, A., Niess, A., Eschweiler, G.W., 2010. Exercise-induced normalization of decreased BDNF serum concentration in elderly women with remitted major depression. *International Journal of Neuropsychopharmacology* 13, 595–602.
- Leal-Cerro, A., Gippini, A., Amaya, M.J., Lage, M., Mato, J.A., Dieguez, C., Casanueva, F.F., 2003. Mechanisms underlying the neuroendocrine response to physical exercise. *Journal of Endocrinological Investigation* 26, 879–885.
- Lee, A.K., Mojtahed-Jaberi, M., Kyriakou, T., Astarloa, E.A., Arno, M., Marshall, N.J., Brain, S.D., O'Dell, S.D., 2010. Effect of high-fat feeding on expression of genes controlling availability of dopamine in mouse hypothalamus. *Nutrition* 26, 411–422.
- Lee, B.H., Kim, Y.K., 2010. The roles of BDNF in the pathophysiology of major depression and in antidepressant treatment. *Psychiatry Investigation* 7, 231–235.
- Leger, D., Guilleminault, C., Dreyfus, J.P., Delahaye, C., Paillard, M., 2000. Prevalence of insomnia in a survey of 12,778 adults in France. *Journal of Sleep Research* 9, 35–42.
- Leonard, B., Maes, M., 2012. Mechanistic explanations how cell-mediated immune activation, inflammation and oxidative and nitrosative stress pathways and their sequels and concomitants play a role in the pathophysiology of unipolar depression. *Neuroscience & Biobehavioral Reviews* 36, 764–785.
- Leproult, R., Copinschi, G., Buxton, O., Van Cauter, E., 1997. Sleep loss results in an elevation of cortisol levels the next evening. *Sleep* 20, 865–870.
- Lin, P.Y., Huang, S.Y., Su, K.P., 2010. A meta-analytic review of polyunsaturated fatty acid compositions in patients with depression. *Biological Psychiatry* 68, 140–147.
- Little, J.P., Safdar, A., Benton, C.R., Wright, D.C., 2011. Skeletal muscle and beyond: the role of exercise as a mediator of systemic mitochondrial biogenesis. *Applied Physiology, Nutrition, and Metabolism* 36, 598–607.
- Liu, W., Xu, Y., Lu, J., Zhang, Y., Sheng, H., Ni, X., 2012a. Swimming exercise ameliorates depression-like behaviors induced by prenatal exposure to glucocorticoids in rats. *Neuroscience Letters* 524, 119–123.
- Liu, Y., Ho, R.C., Mak, A., 2012b. Interleukin (IL)-6, tumour necrosis factor alpha (TNF-alpha) and soluble interleukin-2 receptors (sIL-2R) are elevated in patients with major depressive disorder: a meta-analysis and meta-regression. *Journal of Affective Disorders* 139, 230–239.
- Lopresti, A.L., Hood, S.D., Drummond, P.D., 2012. Multiple antidepressant potential modes of action of curcumin: a review of its anti-inflammatory, monoaminergic, antioxidant, immune-modulating and neuroprotective effects. *Journal of Psychopharmacology* 26, 1512–1524.
- Lucassen, P.J., Meerlo, P., Naylor, A.S., van Dam, A.M., Dayer, A.G., Fuchs, E., Oomen, C.A., Czeh, B., 2010. Regulation of adult neurogenesis by stress, sleep disruption, exercise and inflammation: implications for depression and antidepressant action. *European Neuropsychopharmacology* 20, 1–17.
- Luciano, M., Möttus, R., Starr, J.M., McNeill, G., Jia, X., Craig, L.C., Deary, I.J., 2012. Depressive symptoms and diet: their effects on prospective inflammation levels in the elderly. *Brain, Behavior, and Immunity* 26, 717–720.

- Maes, M., De Ruyter, M., Hobin, P., Suy, E., 1987. Relationship between the dexamethasone suppression test and the L-tryptophan/competing amino acids ratio in depression. *Psychiatric Research* 21, 323–335.
- Maes, M., De Vos, N., Pioli, R., Demedts, P., Wauters, A., Neels, H., Christophe, A., 2000. Lower serum vitamin E concentrations in major depression. Another marker of lowered antioxidant defenses in that illness. *Journal of Affective Disorders* 58, 241–246.
- Maes, M., Galecki, P., Chang, Y.S., Berk, M., 2011a. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 35, 676–692.
- Maes, M., Galecki, P., Verkerk, R., Rief, W., 2011b. Somatization, but not depression, is characterized by disorders in the tryptophan catabolite (TRYCAT) pathway, indicating increased indoleamine 2,3-dioxygenase and lowered kynurenine aminotransferase activity. *Neuroendocrinology Letters* 32, 264–273.
- Maes, M., Leonard, B.E., Myint, A.M., Kubera, M., Verkerk, R., 2011c. The new '5-HT' hypothesis of depression: cell-mediated immune activation induces indoleamine 2,3-dioxygenase, which leads to lower plasma tryptophan and an increased synthesis of detrimental tryptophan catabolites (TRYCATs), both of which contribute to the onset of depression. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 35, 702–721.
- Maes, M., Mihaylova, I., Kubera, M., Uytterhoeven, M., Vrydags, N., Bosmans, E., 2009a. Increased 8-hydroxy-deoxyguanosine, a marker of oxidative damage to DNA, in major depression and myalgic encephalomyelitis/chronic fatigue syndrome. *Neuroendocrinology Letters* 30, 715–722.
- Maes, M., Mihaylova, I., Kubera, M., Uytterhoeven, M., Vrydags, N., Bosmans, E., 2009b. Lower plasma Coenzyme Q10 in depression: a marker for treatment resistance and chronic fatigue in depression and a risk factor for cardiovascular disorder in that illness. *Neuroendocrinology Letters* 30, 462–469.
- Maes, M., Mihaylova, I., Kubera, M., Uytterhoeven, M., Vrydags, N., Bosmans, E., 2010. Increased plasma peroxides and serum oxidized low density lipoprotein antibodies in major depression: markers that further explain the higher incidence of neurodegeneration and coronary artery disease. *Journal of Affective Disorders* 125, 287–294.
- Maes, M., Mihaylova, I., Kubera, M., Uytterhoeven, M., Vrydags, N., Bosmans, E., 2011d. Lower whole blood glutathione peroxidase (GPX) activity in depression, but not in myalgic encephalomyelitis/chronic fatigue syndrome: another pathway that may be associated with coronary artery disease and neuroprogression in depression. *Neuroendocrinology Letters* 32, 133–140.
- Maes, M., Scharpe, S., Meltzer, H.Y., Okayli, G., Bosmans, E., D'Hondt, P., Vanden Bossche, B.V., Cosyns, P., 1994. Increased neopterin and interferon-gamma secretion and lower availability of L-tryptophan in major depression: further evidence for an immune response. *Psychiatric Research* 54, 143–160.
- Maes, M., Van De Vyvere, J., Vandoolaeghe, E., Bril, T.D.P., Wauters, A., Neels, H., 1996. Alterations in iron metabolism and the erythron in major depression: further evidence for a chronic inflammatory process. *Journal of Affective Disorders* 40, 23–33.
- Maes, M., Yirmiya, R., Norberg, J., Brene, S., Hibbeln, J., Perini, G., Kubera, M., Bob, P., Lerer, B., Maj, M., 2009c. The inflammatory & neurodegenerative (I&ND) hypothesis of depression: leads for future research and new drug developments in depression. *Metabolic Brain Disease* 24, 27–53.
- Magalhaes, P.V., Dean, O.M., Bush, A.L., Copolov, D.L., Malhi, G.S., Kohlmann, K., Jeavons, S., Schapkaiz, I., Anderson-Hunt, M., Berk, M., 2011. N-acetylcysteine for major depressive episodes in bipolar disorder. *Revista Brasileira de Psiquiatria* 33, 374–378.
- Maioli, S., Puerta, E., Merino-Serrais, P., Fusari, L., Gil-Bea, F., Rimondini, R., Cedazo-Minguez, A., 2012. Combination of Apolipoprotein E4 and high carbohydrate diet reduces hippocampal BDNF and arc levels and impairs memory in young mice. *Journal of Alzheimer's Disease* 32, 341–355.
- Maletic, V., Robinson, M., Oakes, T., Iyengar, S., Ball, S.G., Russell, J., 2007. Neurobiology of depression: an integrated view of key findings. *International Journal of Clinical Practice* 61, 2030–2040.
- Mamplakou, E., Bountziouka, V., Psaltopoulou, T., Zaimbekis, A., Tsakoundakis, N., Papaerakleous, N., Gotsis, E., Metallinos, G., Pounis, G., Polychronopoulos, E., Lionis, C., Panagiotakos, D., 2010. Urban environment, physical inactivity and unhealthy dietary habits correlate to depression among elderly living in eastern Mediterranean islands: the MEDIS (MEDiterranean Islands Elderly) study. *Journal of Nutrition Health and Aging* 14, 449–455.
- Manber, R., Edinger, J.D., Gress, J.L., San Pedro-Salcedo, M.G., Kuo, T.F., Kalista, T., 2008. Cognitive behavioral therapy for insomnia enhances depression outcome in patients with comorbid major depressive disorder and insomnia. *Sleep* 31, 489–495.
- Manji, H.K., Drevets, W.C., Charney, D.S., 2001. The cellular neurobiology of depression. *Nature Medicine* 7, 541–547.
- Markus, C.R., 2008. Dietary amino acids and brain serotonin function; implications for stress-related affective changes. *NeuroMolecular Medicine* 10, 247–258.
- Martin, A., Prior, R., Shukitt-Hale, B., Cao, G., Joseph, J.A., 2000. Effect of fruits, vegetables, or vitamin E-rich diet on vitamins E and C distribution in peripheral and brain tissues: implications for brain function. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 55, B144–B151.
- Martinowich, K., Lu, B., 2008. Interaction between BDNF and serotonin: role in mood disorders. *Neuropsychopharmacology* 33, 73–83.
- Martinowich, K., Manji, H., Lu, B., 2007. New insights into BDNF function in depression and anxiety. *Nature Neuroscience* 10 (9), 1089–1093.
- Mastorakos, G., Pavlatou, M., Diamanti-Kandarakis, E., Chrousos, G.P., 2005. Exercise and the stress system. *Hormones* 4, 73–89.
- Mathur, N., Pedersen, B.K., 2008. Exercise as a mean to control low-grade systemic inflammation. *Mediators of Inflammation* 2008, 109502.
- Matsuoka, Y., Nishi, D., Yonemoto, N., Hamazaki, K., Hamazaki, T., Hashimoto, K., 2011. Potential role of brain-derived neurotrophic factor in omega-3 fatty acid supplementation to prevent posttraumatic distress after accidental injury: an open-label pilot study. *Psychotherapy and Psychosomatics* 80, 310–312.
- Matsuzawa-Nagata, N., Takamura, T., Ando, H., Nakamura, S., Kurita, S., Misu, H., Ota, T., Yokoyama, M., Honda, M., Miyamoto, K., Kaneko, S., 2008. Increased oxidative stress precedes the onset of high-fat diet-induced insulin resistance and obesity. *Metabolism* 57, 1071–1077.
- Matthes, S., Mosienko, V., Bashammakh, S., Alenina, N., Bader, M., 2010. Tryptophan hydroxylase as novel target for the treatment of depressive disorders. *Pharmacology* 85, 95–109.
- McBride, H.M., Neuspiel, M., Wasiaik, S., 2006. Mitochondria: more than just a powerhouse. *Current Biology* 6, R551–R560.
- McDermott, C.M., LaHoste, G.J., Chen, C., Musto, A., Bazan, N.G., Magee, J.C., 2003. Sleep deprivation causes behavioral, synaptic, and membrane excitability alterations in hippocampal neurons. *Journal of Neuroscience* 23, 9687–9695.
- McDonald, N.C., 2007. Active transportation to school: trends among US school children, 1969–2001. *American Journal of Preventive Medicine* 32, 509–516.
- McNicholas, W.T., 2009. Obstructive sleep apnea and inflammation. *Progress in Cardiovascular Diseases* 51, 392–399.
- Meerlo, P., Koehl, M., van der Borgh, K., Turek, F.W., 2002. Sleep restriction alters the hypothalamic-pituitary-adrenal response to stress. *Journal of Neuroendocrinology* 14, 397–402.
- Meerlo, P., Mistlberger, R.E., Jacobs, B.L., Heller, H.C., McGinty, D., 2009. New neurons in the adult brain: the role of sleep and consequences of sleep loss. *Sleep Medicine Reviews* 13, 187–194.
- Meerlo, P., Sgoifo, A., Suchecki, D., 2008. Restricted and disrupted sleep: effects on autonomic function, neuroendocrine stress systems and stress reactivity. *Sleep Medicine Reviews* 12, 197–210.
- Melancon, M.O., Lorrain, D., Dionne, I.J., 2012. Exercise increases tryptophan availability to the brain in older men age 57–70 years. *Medicine and Science in Sports and Exercise* 44, 881–887.
- Merete, C., Falcon, L.M., Tucker, K.L., 2008. Vitamin B6 is associated with depressive symptomatology in Massachusetts elders. *Journal of the American College of Nutrition* 27, 421–427.
- Meyer, J.H., Ginovart, N., Boovariwala, A., Sagrati, S., Hussey, D., Garcia, A., Young, T., Praschak-Rieder, N., Wilson, A.A., Houle, S., 2006. Elevated monoamine oxidase levels in the brain: an explanation for the monoamine imbalance of major depression. *Archives of General Psychiatry* 63, 1209–1216.
- Michaeli, B., Berger, M.M., Revely, J.P., Tappy, L., Chiolero, R., 2007. Effects of fish oil on the neuro-endocrine responses to an endotoxin challenge in healthy volunteers. *Clinical Nutrition* 26, 70–77.
- Milaneschi, Y., Bandinelli, S., Penninx, B.W., Vogelzangs, N., Corsi, A.M., Lauretani, F., Kisiailiou, A., Vazzana, R., Terracciano, A., Guralnik, J.M., Ferrucci, L., 2011. Depressive symptoms and inflammation increase in a prospective study of older adults: a protective effect of a healthy (Mediterranean-style) diet. *Molecular Psychiatry* 16, 589–590.
- Miller, A.L., 2008. The methylation, neurotransmitter, and antioxidant connections between folate and depression. *Alternative Medicine Review* 13, 216–226.
- Mokhber, N., Namjoo, M., Tara, F., Boskabadi, H., Rayman, M.P., Ghayour-Mobarhan, M., Sahebkar, A., Majidi, M.R., Tavallaie, S., Azimi-Nezhad, M., Shakeri, M.T., Nemati, M., Oladi, M., Mohammadi, M., Ferns, G., 2011. Effect of supplementation with selenium on postpartum depression: a randomized double-blind placebo-controlled trial. *Journal of Maternal-Fetal and Neonatal Medicine* 24, 104–108.
- Moorthy, D., Peter, I., Scott, T.M., Parnell, L.D., Lai, C.Q., Crott, J.W., Ordovas, J.M., Selhub, J., Griffith, J., Rosenberg, I.H., Tucker, K.L., Troen, A.M., 2012. Status of vitamins B-12 and B-6 but not of folate, homocysteine, and the methylenetetrahydrofolate reductase C677T polymorphism are associated with impaired cognition and depression in adults. *Journal of Nutrition* 142, 1554–1560.
- Morrell, M.J., McRobbie, D.W., Quest, R.A., Cummin, A.R., Ghiassi, R., Corfield, D.R., 2003. Changes in brain morphology associated with obstructive sleep apnea. *Sleep Medicine* 4, 451–454.
- Morris, D.W., Trivedi, M.H., Rush, A.J., 2008. Folate and unipolar depression. *Journal of Alternative and Complementary Medicine* 14, 277–285.
- Motivala, S.J., 2011. Sleep and inflammation: psychoneuroimmunology in the context of cardiovascular disease. *Annals of Behavioral Medicine* 42, 141–152.
- Motivala, S.J., Levin, M.J., Oxman, M.N., Irwin, M.R., 2006. Impairments in health functioning and sleep quality in older adults with a history of depression. *Journal of the American Geriatrics Society* 54, 1184–1191.
- Motivala, S.J., Sarfatti, A., Olmos, L., Irwin, M.R., 2005. Inflammatory markers and sleep disturbance in major depression. *Psychosomatic Medicine* 67, 187–194.
- Myint, A.M., Kim, Y.K., Verkerk, R., Scharpe, S., Steinbusch, H., Leonard, B., 2007. Kynurenine pathway in major depression: evidence of impaired neuroprotection. *Journal of Affective Disorders* 98, 143–151.
- Myint, A.M., Leonard, B.E., Steinbusch, H.W., Kim, Y.K., 2005. Th1, Th2, and Th3 cytokine alterations in major depression. *Journal of Affective Disorders* 88, 167–173.
- Nabkasorn, C., Miyai, N., Sootmongkol, A., Junprasert, S., Yamamoto, H., Arita, M., Miyashita, K., 2006. Effects of physical exercise on depression, neuroendocrine stress hormones and physiological fitness in adolescent females with depressive symptoms. *European Journal of Public Health* 16, 179–184.

- Nagahara, A.H., Tuszynski, M.H., 2011. Potential therapeutic uses of BDNF in neurological and psychiatric disorders. *Nature Reviews Drug Discovery* 10, 209–219.
- Nemeroff, C.B., Krishnan, K.R., Reed, D., Leder, R., Beam, C., Dunnick, N.R., 1992. Adrenal gland enlargement in major depression. A computed tomographic study. *Archives of General Psychiatry* 49, 384–387.
- Neylan, T.C., Mueller, S.G., Wang, Z., Metzler, T.J., Lenoci, M., Truran, D., Marmar, C.R., Weiner, M.W., Schuff, N., 2010. Insomnia severity is associated with a decreased volume of the CA3/dentate gyrus hippocampal subfield. *Biological Psychiatry* 68, 494–496.
- Novati, A., Hulshof, H.J., Granic, I., Meerlo, P., 2012. Chronic partial sleep deprivation reduces brain sensitivity to glutamate N-methyl-D-aspartate receptor-mediated neurotoxicity. *Journal of Sleep Research* 21, 3–9.
- Novati, A., Hulshof, H.J., Koolhaas, J.M., Lucassen, P.J., Meerlo, P., 2011. Chronic sleep restriction causes a decrease in hippocampal volume in adolescent rats, which is not explained by changes in glucocorticoid levels or neurogenesis. *Neuroscience* 190, 145–155.
- Novati, A., Roman, V., Cetin, T., Hagewoud, R., den Boer, J.A., Luiten, P.G., Meerlo, P., 2008. Chronically restricted sleep leads to depression-like changes in neurotransmitter receptor sensitivity and neuroendocrine stress reactivity in rats. *Sleep* 31, 1579–1585.
- Ohayon, M.M., 2002. Epidemiology of insomnia: what we know and what we still need to learn. *Sleep Medicine Reviews* 6, 97–111.
- Olson, A.K., Eadie, B.D., Ernst, C., Christie, B.R., 2006. Environmental enrichment and voluntary exercise massively increase neurogenesis in the adult hippocampus via dissociable pathways. *Hippocampus* 16, 250–260.
- Opp, M.R., 2005. Cytokines and sleep. *Sleep Medicine Reviews* 9, 355–364.
- Owen, A.J., Batterham, M.J., Probst, Y.C., Grenyer, B.F., Tapsell, L.C., 2005. Low plasma vitamin E levels in major depression: diet or disease? *European Journal of Clinical Nutrition* 59, 304–306.
- Ozcan, M.E., Gulec, M., Ozerol, E., Polat, R., Akyol, O., 2004. Antioxidant enzyme activities and oxidative stress in affective disorders. *International Clinical Psychopharmacology* 19, 89–95.
- Page, M.M., Robb, E.L., Salway, K.D., Stuart, J.A., 2010. Mitochondrial redox metabolism: aging, longevity and dietary effects. *Mechanisms of Ageing and Development* 131, 242–252.
- Panchal, S.K., Poudyal, H., Iyer, A., Nazer, R., Alam, M.A., Diwan, V., Kauter, K., Sernia, C., Campbell, F., Ward, L., Gobe, G., Fenning, A., Brown, L., 2011. High-carbohydrate, high-fat diet-induced metabolic syndrome and cardiovascular remodeling in rats. *Journal of Cardiovascular Pharmacology* 57, 611–624.
- Pariante, C.M., Lightman, S.L., 2008. The HPA axis in major depression: classical theories and new developments. *Trends in Neurosciences* 31, 464–468.
- Pasco, J.A., Jacka, F.N., Williams, L.J., Evans-Cleverdon, M., Brennan, S.L., Kotowicz, M.A., Nicholson, G.C., Ball, M.J., Berk, M., 2012. Dietary selenium and major depression: a nested case-control study. *Complementary Therapies in Medicine* 20, 119–123.
- Pedersen, B.K., Fischer, C.P., 2007. Physiological roles of muscle-derived interleukin-6 in response to exercise. *Current Opinion in Clinical Nutrition and Metabolic Care* 10, 265–271.
- Pedersen, M., Bruunsgaard, H., Weis, N., Hendel, H.W., Andreassen, B.U., Eldrup, E., Dela, F., Pedersen, B.K., 2003. Circulating levels of TNF-alpha and IL-6-relation to truncal fat mass and muscle mass in healthy elderly individuals and in patients with type-2 diabetes. *Mechanisms of Ageing and Development* 124, 495–502.
- Petersson, H., Risérus, U., McMonagle, J., Gulseth, H.L., Tierney, A.C., Morange, S., Helal, O., Shaw, D.I., Ruano, J.A., López-Miranda, J., Kieć-Wilk, B., Gołabek, I., Blaak, E.E., Saris, W.H., Drevon, C.A., Lovegrove, J.A., Roche, H.M., Basu, S., 2010. Effects of dietary fat modification on oxidative stress and inflammatory markers in the LIPGENE study. *British Journal of Nutrition* 104, 1357–1362.
- Piecznik, S.R., Neustadt, J., 2007. Mitochondrial dysfunction and molecular pathways of disease. *Experimental and Molecular Pathology* 83, 84–92.
- Pittaluga, M., Parisi, P., Sabatini, S., Ceci, R., Caporossi, D., Valeria Catani, M., Savini, I., Avigliano, L., 2006. Cellular and biochemical parameters of exercise-induced oxidative stress: relationship with training levels. *Free Radical Research* 40, 607–614.
- Plaisance, E.P., Grandjean, P.W., 2006. Physical activity and high-sensitivity C-reactive protein. *Sports Medicine* 36, 443–458.
- Ploeger, H.E., Takken, T., De Greef, M.H., Timmons, B.W., 2009. The effects of acute and chronic exercise on inflammatory markers in children and adults with a chronic inflammatory disease: a systematic review. *Exercise Immunology Review* 15, 6–41.
- Popkin, B.M., Nielsen, S.J., 2003. The sweetening of the world's diet. *Obesity Research* 11, 1325–1332.
- Pritchett, C.E., Hajnal, A., 2011. Obesogenic diets may differentially alter dopamine control of sucrose and fructose intake in rats. *Physiology & Behavior* 104, 111–116.
- Radak, Z., Chung, H.Y., Goto, S., 2008. Systemic adaptation to oxidative challenge induced by regular exercise. *Free Radical Biology and Medicine* 44, 153–159.
- Raison, C.L., Dantzer, R., Kelley, K.W., Lawson, M.A., Woolwine, B.J., Vogt, G., Spivey, J.R., Saito, K., Miller, A.H., 2010a. CSF concentrations of brain tryptophan and kynurenines during immune stimulation with IFN-alpha: relationship to CNS immune responses and depression. *Molecular Psychiatry* 15, 393–403.
- Raison, C.L., Miller, A.H., 2011. Is depression an inflammatory disorder? *Current Psychiatry Reports* 13, 467–475.
- Raison, C.L., Rye, D.B., Woolwine, B.J., Vogt, G.J., Bautista, B.M., Spivey, J.R., Miller, A.H., 2010b. Chronic interferon-alpha administration disrupts sleep continuity and depth in patients with hepatitis C: association with fatigue, motor slowing, and increased evening cortisol. *Biological Psychiatry* 68, 942–949.
- Ramanathan, L., Gulyani, S., Nienhuis, R., Siegel, J.M., 2002. Sleep deprivation decreases superoxide dismutase activity in rat hippocampus and brainstem. *NeuroReport* 13, 1387–1390.
- Rao, J.S., Ertley, R.N., Lee, H.J., DeMar Jr., J.C., Arnold, J.T., Rapoport, S.I., Bazinet, R.P., 2007. n-3 Polyunsaturated fatty acid deprivation in rats decreases frontal cortex BDNF via a p38 MAPK-dependent mechanism. *Molecular Psychiatry* 12, 36–46.
- Rechtschaffen, A., Gilliland, M.A., Bergmann, B.M., Winter, J.B., 1983. Physiological correlates of prolonged sleep deprivation in rats. *Science* 221, 182–184.
- Reimund, E., 1994. The free radical flux theory of sleep. *Medical Hypotheses* 43, 231–233.
- Ribeiro, M.C., Barbosa, N.B., De Almeida, T.M., Parcianello, L.M., Perottoni, J., De Avila, D.S., Rocha, J.B., 2009. High-fat diet and hydrochlorothiazide increase oxidative stress in brain of rats. *Cell Biochemistry & Function* 27, 473–478.
- Richard, C., Couture, P., Desroches, S., Lamarche, B., 2012. Effect of the Mediterranean diet with and without weight loss on markers of inflammation in men with metabolic syndrome. *Obesity*, <http://dx.doi.org/10.1038/oby.2012.148>, Epub ahead of print.
- Riemann, D., Voderholzer, U., 2003. Primary insomnia: a risk factor to develop depression? *Journal of Affective Disorders* 76, 255–259.
- Riemann, D., Voderholzer, U., Spiegelhalter, K., Hornyak, M., Buysse, D.J., Nissen, C., Hennig, J., Perlis, M.L., Van Elst, L.T., Feige, B., 2007. Chronic insomnia and MRI-measured hippocampal volumes: a pilot study. *Sleep* 30, 955–958.
- Rimer, J., Dwan, K., Lawlor, D.A., Greig, C.A., McMurdo, M., Morley, W., Mead, G.E., 2012. Exercise for depression. *Cochrane Database of Systematic Reviews* 7, CD004366.
- Rodenbeck, A., Hajak, G., 2001. Neuroendocrine dysregulation in primary insomnia. *Revista de Neurología* 157, S57–S61.
- Roman, V., Hagewoud, R., Luiten, P.G., Meerlo, P., 2006. Differential effects of chronic partial sleep deprivation and stress on serotonin-1A and muscarinic acetylcholine receptor sensitivity. *Journal of Sleep Research* 15, 386–394.
- Roman, V., Van der Borght, K., Leemburg, S.A., Van der Zee, E.A., Meerlo, P., 2005a. Sleep restriction by forced activity reduces hippocampal cell proliferation. *Brain Research* 1065, 53–59.
- Roman, V., Walstra, I., Luiten, P.G., Meerlo, P., 2005b. Too little sleep gradually desensitizes the serotonin 1A receptor system. *Sleep* 28, 1505–1510.
- Sable, P.S., Dangat, K.D., Joshi, A.A., Joshi, S.R., 2012. Maternal omega 3 fatty acid supplementation during pregnancy to a micronutrient-imbalanced diet protects postnatal reduction of brain neurotrophins in the rat offspring. *Neuroscience* 217, 46–55.
- Sanchez-Villegas, A., Delgado-Rodriguez, M., Alonso, A., Schlatter, J., Lahortiga, F., Serra Majem, L., Martinez-Gonzalez, M.A., 2009. Association of the Mediterranean dietary pattern with the incidence of depression: the Seguimiento Universidad de Navarra/University of Navarra follow-up (SUN) cohort. *Archives of General Psychiatry* 66, 1090–1098.
- Sanchez-Villegas, A., Galbete, C., Martinez-Gonzalez, M.A., Martinez, J.A., Razquin, C., Salas-Salvado, J., Estruch, R., Buil-Cosiales, P., Marti, A., 2011. The effect of the Mediterranean diet on plasma brain-derived neurotrophic factor (BDNF) levels: the PREDIMED-NAVARRA randomized trial. *Nutritional Neuroscience* 14, 195–201.
- Sanchez-Villegas, A., Toledo, E., De Irala, J., Ruiz-Canela, M., Pla-Vidal, J., Martinez-Gonzalez, M.A., 2012. Fast-food and commercial baked goods consumption and the risk of depression. *Public Health Nutrition* 15, 424–432.
- Santos, R.V., Tufik, S., De Mello, M.T., 2007. Exercise, sleep and cytokines: is there a relation? *Sleep Medicine Reviews* 11, 231–239.
- Sarandol, A., Sarandol, E., Eker, S.S., Erdinc, S., Vatansver, E., Kirli, S., 2007. Major depressive disorder is accompanied with oxidative stress: short-term antidepressant treatment does not alter oxidative-antioxidative systems. *Human Psychopharmacology* 22, 67–73.
- Savard, J., Simard, S., Ivers, H., Morin, C.M., 2005. Randomized study on the efficacy of cognitive-behavioral therapy for insomnia secondary to breast cancer, part I: sleep and psychological effects. *Journal of Clinical Oncology: Official Journal of the American Society of Clinical Oncology* 23, 6083–6096.
- Schwarz, A.J., Brasel, J.A., Hintz, R.L., Mohan, S., Cooper, D.M., 1996. Acute effect of brief low- and high-intensity exercise on circulating insulin-like growth factor (IGF) I, II, and IGF-binding protein-3 and its proteolysis in young healthy men. *Journal of Clinical Endocrinology and Metabolism* 81, 3492–3497.
- Schweiger, U., Broocks, A., Tuschl, R.J., Pirke, K.M., 1989. Serotonin turnover in rat brain during semistarvation with high-protein and high-carbohydrate diets. *Journal of Neural Transmission* 77, 131–139.
- Sen, S., Duman, R., Sanacora, G., 2008. Serum brain-derived neurotrophic factor, depression, and antidepressant medications: meta-analyses and implications. *Biological Psychiatry* 64, 527–532.
- Shao, L., Martin, M.V., Watson, S.J., Schatzberg, A., Akil, H., Myers, R.M., Jones, E.G., Bunney, W.E., Vawter, M.P., 2008. Mitochondrial involvement in psychiatric disorders. *Annals of Medicine* 40, 281–295.
- Skalicky, J., Muzakova, V., Kandar, R., Meloun, M., Rousar, T., 2009. Oxidative stress and metabolic syndrome in obese adults with and without controlled diet restriction. *Bratislava Medical Journal* 110, 152–157.
- Skarupski, K.A., Tangney, C., Li, H., Ouyang, B., Evans, D.A., Morris, M.C., 2010. Longitudinal association of vitamin B-6, folate, and vitamin B-12 with depressive symptoms among older adults over time. *American Journal of Clinical Nutrition* 92, 330–335.
- Sockalingam, S., Abbey, S.E., Alosaimi, F., Novak, M., 2010. A review of sleep disturbance in hepatitis C. *Journal of Clinical Gastroenterology* 44, 38–45.

- Song, M.R., Lee, Y.S., Baek, J.D., Miller, M., 2012. Physical activity status in adults with depression in the National Health and Nutrition Examination Survey, 2005–2006. *Public Health Nursing* 29, 208–217.
- Spiegel, K., Leproult, R., Van Cauter, E., 1999. Impact of sleep debt on metabolic and endocrine function. *Lancet* 354, 1435–1439.
- Stahl, S.M., 2008. L-methylfolate: a vitamin for your monoamines. *Journal of Clinical Psychiatry* 69, 1352–1353.
- Starkie, R., Ostrowski, S.R., Jauffred, S., Febbraio, M., Pedersen, B.K., 2003. Exercise and IL-6 infusion inhibit endotoxin-induced TNF- $\alpha$  production in humans. *FASEB Journal* 17, 884–886.
- Steiner, J., Walter, M., Gos, T., Guillemin, G.J., Bernstein, H.G., Sarnyai, Z., Mawrin, C., Brisch, R., Bielau, H., Meyer zu Schwabedissen, L., Bogerts, B., Myint, A.M., 2011. Severe depression is associated with increased microglial quinolinic acid in subregions of the anterior cingulate gyrus: evidence for an immunomodulated glutamatergic neurotransmission? *Journal of Neuroinflammation* 8, 94.
- Stephens, T., 1988. Physical activity and mental health in the United States and Canada: evidence from four population surveys. *Preventive Medicine* 17, 35–47.
- Stewart, R., Hirani, V., 2012. Relationship between depressive symptoms, anemia, and iron status in older residents from a national survey population. *Psychosomatic Medicine* 74, 208–213.
- Stranahan, A.M., Cutler, R.G., Button, C., Telljohann, R., Mattson, M.P., 2011. Diet-induced elevations in serum cholesterol are associated with alterations in hippocampal lipid metabolism and increased oxidative stress. *Journal of Neurochemistry* 118, 611–615.
- Stranahan, A.M., Lee, K., Mattson, M.P., 2008. Central mechanisms of HPA axis regulation by voluntary exercise. *NeuroMolecular Medicine* 10, 118–127.
- Su, K.P., 2009. Biological mechanism of antidepressant effect of omega-3 fatty acids: how does fish oil act as a 'mind-body interface'? *Neurosignals* 17, 144–152.
- Suarez, E.C., 2008. Self-reported symptoms of sleep disturbance and inflammation, coagulation, insulin resistance and psychosocial distress: evidence for gender disparity. *Brain, Behavior, and Immunity* 22, 960–968.
- Sublette, M.E., Ellis, S.P., Geant, A.L., Mann, J.J., 2011. Meta-analysis of the effects of eicosapentaenoic acid (EPA) in clinical trials in depression. *Journal of Clinical Psychiatry* 72, 1577–1584.
- Suchecki, D., Lobo, L.L., Hipolide, D.C., Tufik, S., 1998. Increased ACTH and corticosterone secretion induced by different methods of paradoxical sleep deprivation. *Journal of Sleep Research* 7, 276–281.
- Suer, C., Dolu, N., Artis, A.S., Sahin, L., Yilmaz, A., Cetin, A., 2011. The effects of long-term sleep deprivation on the long-term potentiation in the dentate gyrus and brain oxidation status in rats. *Neuroscience Research* 70, 71–77.
- Szewczyk, B., Kubera, M., Nowak, G., 2011. The role of zinc in neurodegenerative inflammatory pathways in depression. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 35, 693–701.
- Szewczyk, B., Poleszak, E., Wlaz, P., Wrobel, A., Blicharska, E., Cichy, A., Dybala, M., Siwek, A., Pomierny-Chamiolo, L., Piotrowska, A., Branski, P., Pilc, A., Nowak, G., 2009. The involvement of serotonergic system in the antidepressant effect of zinc in the forced swim test. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 33, 323–329.
- Taafe, D.R., Harris, T.B., Ferrucci, L., Rowe, J., Seeman, T.E., 2000. Cross-sectional and prospective relationships of interleukin-6 and C-reactive protein with physical performance in elderly persons: MacArthur studies of successful aging. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 55, M709–M715.
- Taylor, D.J., Lichstein, K.L., Durrence, H.H., Reid, B.W., Bush, A.J., 2005. Epidemiology of insomnia, depression, and anxiety. *Sleep* 28, 1457–1464.
- Teegarden, S.L., Scott, A.N., Bale, T.L., 2009. Early life exposure to a high fat diet promotes long-term changes in dietary preferences and central reward signaling. *Neuroscience* 162, 924–932.
- Teychenne, M., Ball, K., Salmon, J., 2010. Sedentary behavior and depression among adults: a review. *International Journal of Behavioral Medicine* 17, 246–254.
- Thamaraiselvi, K., Mathangi, D.C., Subhashini, A.S., 2012. Effect of increase in duration of REM sleep deprivation on lipid peroxidation. *International Journal of Biological and Medical Research* 3, 1754–1759.
- Thomas, N.E., Williams, D.R., 2008. Inflammatory factors, physical activity, and physical fitness in young people. *Scandinavian Journal of Medicine and Science in Sports* 18, 543–556.
- Toker, L., Amar, S., Bersudsky, Y., Benjamin, J., Klein, E., Agam, G., 2010. The biology of tryptophan depletion and mood disorders. *Israel Journal of Psychiatry & Related Sciences* 47, 46–55.
- Traustadottir, T., Bosch, P.R., Cantu, T., Matt, K.S., 2004. Hypothalamic-pituitary-adrenal axis response and recovery from high-intensity exercise in women: effects of aging and fitness. *Journal of Clinical Endocrinology & Metabolism* 89, 3248–3254.
- Urpi-Sarda, M., Casas, R., Chiva-Blanch, G., Romero-Mamani, E.S., Valderas-Martínez, P., Salas-Salvadó, J., Covas, M.I., Toledo, E., Andres-Lacueva, C., Llorach, R., García-Arellano, A., Bulló, M., Ruiz-Gutiérrez, V., Lamuela-Raventós, R.M., Estruch, R., 2012. The Mediterranean diet pattern and its main components are associated with lower plasma concentrations of tumor necrosis factor receptor 60 in patients at high risk for cardiovascular disease. *Journal of Nutrition* 142, 1019–1025.
- Vahdat Shariatpanaahi, M., Vahdat Shariatpanaahi, Z., Moshtaaghi, M., Shahbaazi, S.H., Abadi, A., 2007. The relationship between depression and serum ferritin level. *European Journal of Clinical Nutrition* 61, 532–535.
- Vallance, J.K., Winkler, E.A., Gardiner, P.A., Healy, G.N., Lynch, B.M., Owen, N., 2011. Associations of objectively-assessed physical activity and sedentary time with depression: NHANES (2005–2006). *Preventive Medicine* 53, 284–288.
- van Leeuwen, W.M., Lehto, M., Karisola, P., Lindholm, H., Luukkainen, R., Sallinen, M., Harma, M., Porkka-Heiskanen, T., Alenius, H., 2009. Sleep restriction increases the risk of developing cardiovascular diseases by augmenting proinflammatory responses through IL-17 and CRP. *PLoS One* 4, e4589.
- Vgontzas, A.N., Bixler, E.O., Lin, H.M., Prolo, P., Mastorakos, G., Vela-Bueno, A., Kales, A., Chrousos, G.P., 2001. Chronic insomnia is associated with nocturnal activation of the hypothalamic-pituitary-adrenal axis: clinical implications. *Journal of Clinical Endocrinology & Metabolism* 86, 3787–3794.
- Vgontzas, A.N., Papanicolaou, D.A., Bixler, E.O., Litsikas, A., Zachman, K., Kales, A., Prolo, P., Wong, M.L., Licinio, J., Gold, P.W., Hermida, R.C., Mastorakos, G., Chrousos, G.P., 1999. Circadian interleukin-6 secretion and quantity and depth of sleep. *Journal of Clinical Endocrinology & Metabolism* 84, 2603–2607.
- Vgontzas, A.N., Tsigos, C., Bixler, E.O., Stratakis, C.A., Zachman, K., Kales, A., Vela-Bueno, A., Chrousos, G.P., 1998. Chronic insomnia and activity of the stress system: a preliminary study. *Journal of Psychosomatic Research* 45, 21–31.
- Vgontzas, A.N., Zoumakis, E., Bixler, E.O., Lin, H.M., Follett, H., Kales, A., Chrousos, G.P., 2004. Adverse effects of modest sleep restriction on sleepiness, performance, and inflammatory cytokines. *Journal of Clinical Endocrinology & Metabolism* 89, 2119–2126.
- Vgontzas, A.N., Zoumakis, M., Papanicolaou, D.A., Bixler, E.O., Prolo, P., Lin, H.M., Vela-Bueno, A., Kales, A., Chrousos, G.P., 2002. Chronic insomnia is associated with a shift of interleukin-6 and tumor necrosis factor secretion from nighttime to daytime. *Metabolism* 51, 887–892.
- Vina, J., Gomez-Cabrera, M.C., Borras, C., Froio, T., Sanchis-Gomar, F., Martinez-Bello, V.E., Pallardo, F.V., 2009. Mitochondrial biogenesis in exercise and in ageing. *Advanced Drug Delivery Reviews* 61, 1369–1374.
- Vitetta, L., Anton, B., 2007. Lifestyle and nutrition, caloric restriction, mitochondrial health and hormones: scientific interventions for anti-aging. *Clinical Interventions in Aging* 2, 537–543.
- Vivar, C., Potter, M.C., van Praag, H., 2012. All about running: synaptic plasticity, growth factors and adult hippocampal neurogenesis. *Current Topics in Behavioral Neurosciences*, [http://dx.doi.org/10.1007/7854\\_2012\\_220](http://dx.doi.org/10.1007/7854_2012_220), Epub ahead of print.
- Vollaard, N.B., Shearman, J.P., Cooper, C.E., 2005. Exercise-induced oxidative stress: myths, realities and physiological relevance. *Sports Medicine* 35, 1045–1062.
- Vucetic, Z., Carlin, J.L., Totoki, K., Reyes, T.M., 2012. Epigenetic dysregulation of the dopamine system in diet-induced obesity. *Journal of Neurochemistry* 120, 891–898.
- Vuckovic, M.G., Li, Q., Fisher, B., Nacca, A., Leahy, R.M., Walsh, J.P., Mukherjee, J., Williams, C., Jakowec, M.W., Petzinger, G.M., 2010. Exercise elevates dopamine D2 receptor in a mouse model of Parkinson's disease: in vivo imaging with [(1)(8)F]fallypride. *Movement Disorders* 25, 2777–2784.
- Wang, G.J., Volkow, N.D., Fowler, J.S., Franceschi, D., Logan, J., Pappas, N.R., Wong, C.T., Netusil, N., 2000. PET studies of the effects of aerobic exercise on human striatal dopamine release. *Journal of Nuclear Medicine* 41, 1352–1356.
- Wang, Y.C., Bleich, S.N., Gortmaker, S.L., 2008. Increasing caloric contribution from sugar-sweetened beverages and 100% fruit juices among US children and adolescents, 1988–2004. *Pediatrics* 121, e1604–e1614.
- Wei, Y.C., Zhou, F.L., He, D.L., Bai, J.R., Hui, L.Y., Wang, X.Y., Nan, K.J., 2009. The level of oxidative stress and the expression of genes involved in DNA-damage signaling pathways in depressive patients with colorectal carcinoma. *Journal of Psychosomatic Research* 66, 259–266.
- Weicker, H., Struder, H.K., 2001. Influence of exercise on serotonergic neuromodulation in the brain. *Amino Acids* 20, 35–47.
- Wichers, M.C., Koek, G.H., Robaey, G., Verkerk, R., Scharpe, S., Maes, M., 2005. IDO and interferon- $\alpha$ -induced depressive symptoms: a shift in hypothesis from tryptophan depletion to neurotoxicity. *Molecular Psychiatry* 10, 538–544.
- Williams, A.L., Cotter, A., Sabina, A., Girard, C., Goodman, J., Katz, D.L., 2005. The role for vitamin B-6 as treatment for depression: a systematic review. *Family Practice* 22, 532–537.
- Wipfli, B., Landers, D., Nagoshi, C., Ringenbach, S., 2011. An examination of serotonin and psychological variables in the relationship between exercise and mental health. *Scandinavian Journal of Medicine and Science in Sports* 21, 474–481.
- Wu, A., Ying, Z., Gomez-Pinilla, F., 2004. Dietary omega-3 fatty acids normalize BDNF levels, reduce oxidative damage, and counteract learning disability after traumatic brain injury in rats. *Journal of Neurotrauma* 21, 1457–1467.
- Wu, C.W., Chang, Y.T., Yu, L., Chen, H.L., Jen, C.J., Wu, S.Y., Lo, C.P., Kuo, Y.M., 2008. Exercise enhances the proliferation of neural stem cells and neurite growth and survival of neuronal progenitor cells in dentate gyrus of middle-aged mice. *Journal of Applied Physiology* 105, 1585–1594.
- Wurtman, R.J., Wurtman, J.J., 1995. Brain serotonin, carbohydrate-craving, obesity and depression. *Obesity Research* 3 (4), 477S–480S.
- Yamada-Goto, N., Katsuura, G., Ochi, Y., Ebihara, K., Kusakabe, T., Hosoda, K., Nakao, K., 2012. Impairment of fear-conditioning responses and changes of brain neurotrophic factors in diet-induced obese mice. *Journal of Neuroendocrinology* 24, 1120–1125.
- Yi, S.A.N., Poudel-Tandukar, K., Nonaka, D., Matsushita, Y., Hori, A., Mizoue, T., 2011. Association between serum ferritin concentrations and depressive symptoms in Japanese municipal employees. *Psychiatry Research* 189, 368–372.

- Yook, K., Lee, S.H., Ryu, M., Kim, K.H., Choi, T.K., Suh, S.Y., Kim, Y.W., Kim, B., Kim, M.Y., Kim, M.J., 2008. Usefulness of mindfulness-based cognitive therapy for treating insomnia in patients with anxiety disorders: a pilot study. *Journal of Nervous and Mental Disease* 196, 501–503.
- Yubero-Serrano, E.M., Gonzalez-Guardia, L., Rangel-Zuñiga, O., Delgado-Casado, N., Delgado-Lista, J., Perez-Martinez, P., Garcia-Rios, A., Caballero, J., Marin, C., Gutierrez-Mariscal, F.M., Tinahones, F.J., Villalba, J.M., Tunes, I., Perez-Jimenez, F., Lopez-Miranda, J., 2011. Postprandial antioxidant gene expression is modified by Mediterranean diet supplemented with coenzyme Q(10) in elderly men and women. *Age*. <http://dx.doi.org/10.1007/s11357-011-9331-4>, Epub ahead of print.
- Yubero-Serrano, E.M., Gonzalez-Guardia, L., Rangel-Zuniga, O., Delgado-Lista, J., Gutierrez-Mariscal, F.M., Perez-Martinez, P., Delgado-Casado, N., Cruz-Teno, C., Tinahones, F.J., Villalba, J.M., Perez-Jimenez, F., Lopez-Miranda, J., 2012. Mediterranean diet supplemented with coenzyme Q10 modifies the expression of proinflammatory and endoplasmic reticulum stress-related genes in elderly men and women. *Journals of Gerontology Series A: Biological Sciences and Medical Sciences* 67, 3–10.