



Contents lists available at [ScienceDirect](#)

## Journal of Affective Disorders

journal homepage: [www.elsevier.com/locate/jad](http://www.elsevier.com/locate/jad)



### Corrigendum to “Omega-6 fatty acids and greater likelihood of suicide risk and major depression in early pregnancy” [*J. Affect. Disord.* 152–154 (2014) 76–82]



Vaz J.S., Kac G., Nardi A.E., Hibbeln J.R.

*Nutritional Epidemiology Observatory, Josué de Castro Nutrition Institute, Rio de Janeiro Federal University, Rio de Janeiro, Brazil*

#### 2. Methods

##### 2.2. Psychiatric assessment

SR was assessed by means the “suicidality” level obtained from the MINI interview. In this module, six questions were asked, and positive answers received a unique score, as follows: “In the past month did you: (Christian, 2012) think that you would be better off dead or wish you were dead? (Score 1); (Hibbeln and Davis, 2009) want to harm yourself or to hurt or to injure yourself? (Score 2); (Hibbeln, 2009) think about suicide? (Score 6); (Milte et al., 2009) have a suicide plan? (Score 10); (Sublette et al., 2011) attempt suicide? (Score 10); (Lin et al., 2010) during your past, did you ever make a suicide attempt? (Score 4).

#### **Corrigendum:**

SR was assessed by means the “suicidality” level obtained from the MINI interview (version 5.0.0). In this module, six questions were asked, and positive answers received a unique score, as follows: “In the past month did you (1) think that you would be better off dead or wish you were dead? (Score 1); (2) want to harm yourself or to hurt or to injure yourself? (Score 2); (3) think about suicide? (Score 6); (4) have a suicide plan? (Score 10); (5) attempt suicide? (Score 10); and (6) during your past, did you ever make a suicide attempt? (Score 4)”.

DOI of original article: <http://dx.doi.org/10.1016/j.jad.2013.04.045>

<http://dx.doi.org/10.1016/j.jad.2015.03.014>

0165-0327/© 2015 Elsevier B.V. All rights reserved.