

Challenging the effectiveness of green tea in primary and tertiary cancer prevention

Hirota Fujiki · Kazue Imai · Kei Nakachi ·
Masahito Shimizu · Hisataka Moriwaki ·
Masami Suganuma

Received: 11 April 2012 / Accepted: 22 May 2012 / Published online: 15 June 2012
© Springer-Verlag 2012

Abstract

Purpose Drinking green tea daily is part of Japanese culture, and various studies have revealed that green tea is a cancer preventive. We here review our progress in cancer prevention with green tea on 12 main topics, from basic to clinical level.

Topics and methods Biochemical and biological studies of green tea catechins, a prospective cohort study, preclinical safety trials with tablets of green tea extract, double-blind randomized clinical phase II prevention trial for recurrence of colorectal adenomas, and synergistically enhanced inhibition by the combination of green tea catechins and anticancer drugs. All results were significant, including human studies with informed consent.

Results Drinking 10 Japanese-size cups of green tea per day delayed the cancer onset of humans 7 years for females. For tertiary cancer prevention, consuming 10 cups of green tea per day fortified by green tea tablets, 50 %, significantly prevented the recurrence of colorectal

adenomas. A minimum effective amount of green tea catechins for cancer prevention was found in humans. In addition, the combination of green tea catechins and anti-cancer drugs engendered a new cancer therapeutic strategy. **Conclusion** The consumption of 10 Japanese-size cups of green tea per day is a significant factor in primary cancer prevention for the general population, and the preventive effect on recurrence of colorectal adenomas in patients is vital evidence in tertiary cancer prevention.

Keywords Apoptosis · Delayed cancer onset · GADD153 · Green tea tablet · Phase II prevention trial · Prospective cohort study · TNF- α

Introduction

Japanese have the longest life span in the world, 86 years for females and 79 years for males. Even so, cancer mortality increases after age 60 and is the highest for age groups around 80 years of age for both males and females (Fig. 1). The multistage carcinogenesis study of B. Vogelstein revealed that the first neoplastic change in cells in human colon development starts many years earlier, probably 20–30 years earlier than the clinical appearance of cancer in patients (Vogelstein et al. 1988). M. Sporn coined the term “Cancer chemoprevention,” and defined it as “prevention of the occurrence of cancer by administration of one or several compounds (Sporn et al. 1976).” Today, it is generally accepted that it is possible to delay the clinical appearance of cancer by slowing down the development of cancer. At present, for example, 50 % of breast cancer of high-risk group can be prevented by the administration of tamoxifen, in the United States (Fisher et al. 1998).

H. Fujiki (✉)
Faculty of Pharmaceutical Sciences, Tokushima Bunri
University, Tokushima 770-8514, Japan
e-mail: uv4h-fjk@asahi-net.or.jp

K. Imai · K. Nakachi
Department of Radiobiology and Molecular Epidemiology,
Radiation Effects Research Foundation,
Hiroshima 732-0815, Japan

M. Shimizu · H. Moriwaki
Department of Medicine, Gifu University Graduate School
of Medicine, Gifu 501-1194, Japan

M. Suganuma
Research Institute for Clinical Oncology,
Saitama Cancer Center, Saitama 362-0806, Japan




Fig. 1 Cancer mortality by age in Japan (1997). *M* male, *F* female

When research in cancer chemoprevention began in Japan in 1983, we were lucky to receive tannins or polyphenols isolated from medicinal plants and herbs from T. Okuda at Okayama University (Okuda et al. 1985). To screen for cancer preventive agents, we tried to find inhibitors of tumor promotion in two-stage carcinogenesis experiments on mouse skin, consisting of initiation and tumor promotion. The three diterpene ester tumor promoters discovered by E. Hecker were very useful for our further study (Hecker et al. 1984). From a total of 30 polyphenols, (–)-epigallocatechin gallate (EGCG) and penta-*O*-galloyl- β -D-glucose (5GG) showed binding to the phorbol ester receptor and inhibited 12-*O*-tetradecanoylphorbol-13-acetate (TPA)-induced activation of protein kinase C (PKC). In 1987, we first reported, in the British journal *Phytotherapy Research*, that EGCG prevented tumor promotion of teleocidin, one of the TPA-type tumor promoters on mouse skin (Yoshizawa et al. 1987). Later, we also confirmed that repeated applications of 5 mg EGCG completely prevented tumor promotion of okadaic acid (Yoshizawa et al. 1992), a potent tumor promoter working through the inhibition of protein phosphatases 1 and 2A. The tumor-promoting activity by okadaic acid is as potent as TPA (Suganuma et al. 1988).

After we published the results in 1987, the Journal “New Scientist” soon introduced our results with EGCG in an article entitled “Green tea cuts cancerous growths” (November 12, 1987). In winter of 1987, an Australian TV scientific series entitled “Beyond 2000” visited us at the National Cancer Center Research Institute in Tokyo to make a TV film about our research. This 8-min-long TV film was shown all over the world in 1988 and 1989, and many overseas scientists took greater interest in cancer prevention with green tea than Japanese scientists. A putative carcinogenic potential for tannins or polyphenols suspected previously was eliminated by establishing the non-mutagenicity of EGCG and green tea catechins (Okuda et al. 1984), and by EGCG inhibition of TPA-induced activation of PKC (Yoshizawa et al. 1987).

Moreover, EGCG and green tea extract are non-toxic for rodents and humans (Japanese). These findings encouraged us to intensively study cancer prevention by green tea (Fujiki and Suganuma 2002; Fujiki 2005).

Results and discussion

Green tea and green tea extract

Green tea is made from steamed fresh tea leaves and is a non-oxidized non-fermented product containing at least four green tea catechins—EGCG, (–)-epigallocatechin (EGC), (–)-epicatechin (EC) and (–)-epicatechin gallate (ECG)—and caffeine on high-performance liquid chromatography (HPLC). EGCG is the main constituent (Fujiki and Okuda 1992). In addition to EGCG, we used for our experiments green tea extract, the dried green tea infusion containing all green tea catechins, a condensed form of green tea, and Japanese have been drinking green tea for 800 years. Green tea catechins are active in preventing carcinogenesis in a wide range of target organs, including the digestive tract, lung, liver, pancreas, breast, bladder, prostate and skin in rodents (Fujita et al. 1989; Yamane et al. 1991, 1995; Conney et al. 1992; Wang et al. 1992; Narisawa and Fukaura 1993; Yang and Wang 1993; Nishida et al. 1994; NCI et al. 1996; Fujiki et al. 1996; Gupta et al. 2001). Moreover, drinking 0.05 and 0.1 % EGCG solutions significantly prevented the spontaneous metastasis of B16-BL6 cells from foot pad to the lungs of male C57BL/6 mice (Taniguchi et al. 1992). It is interesting to note that green tea catechins are active in most of these organs.

Tissue distribution of ^3H -EGCG

To prove the systemic effects of EGCG, ^3H -EGCG was intubated into the stomach of mice. The microautoradiography of the lungs showed silver grains of radioactive EGCG in some cells, but not all (Suganuma et al. 1998), apparent confirmation that EGCG was incorporated from the digestive tract into the lungs, one of the target organs. Microautoradiography of human lung cancer cell line PC-9 cells treated with ^3H -EGCG showed silver grains in the membrane, cytosol and nuclei, and that ^3H -EGCG had been incorporated from culture medium into the cells (Okabe et al. 1997). Using cold spray ionization-mass spectrometry and surface plasmon resonance assay (Biacore), the direct binding of EGCG to single-strand 18 mers of DNA and RNA was demonstrated (Kuzuhara et al. 2006). The incorporation of ^3H -EGCG into target organs is shown in Table 1. Various amounts of the total administered radioactivity were found in the digestive tract, liver, brain, kidney, lung, pancreas and skin 24 h after intubation

Table 1 Incorporation of ³H-EGCG into target organs

Organs	% of total administered radioactivity (24 h after)	Reduction in tumor incidence
Stomach	3.93	62.0 → 31.0
Duodenum	0.35	63.0 → 20.0
Small intestine	5.69	ND
Colon	4.52	77.3 → 38.1 67.0 → 33.0 ^a
Liver	0.89	83.3 → 52.2
Brain	0.32	ND
Kidney	0.28	ND
Lung	0.16	96.3 → 65.5
Pancreas	0.07	54.0 → 28.0
Skin	1.9 × 10 ⁴ /100 mg ^b	65.0 → 28.0 ^a

ND, not determined

^a Green tea extract

^b dpm

(Suganuma et al. 1998). The right column shows the reduction in tumor incidence in carcinogenesis by EGCG and green tea extract in rodents (Table 1) (Fujiki et al. 1996). Thus, radioactivity was found in the organs where EGCG and green tea extract had previously been shown to inhibit carcinogenesis.

Significance in multiple administrations of EGCG

Japanese drink green tea throughout the day. To study the effects of frequent consumption, ³H-EGCG was intubated into the stomach of mice at 6-h intervals. Duplicate administrations enhanced incorporation of ³H-EGCG 4- to 9-fold in most organs compared with a single administration, suggesting the accumulation of EGCG in cells. Radioactivity in blood and urine increased as well. We named this synergistic enhancement by EGCG the “Fujiki-Suganuma Effect,” which is supported by frequent drinking of green tea (Table 2) (Suganuma et al. 1998). To further study the effects on enhanced incorporation of EGCG by multiple administrations, the induction of *growth arrest and DNA damage-inducible 153 (GADD153)* and *p21* gene expressions in human lung cancer cell line A549 cells was determined at 6-h intervals. *GADD153 (CHOP)* is an apoptosis-regulating gene. The overexpression of *GADD153* gene induces apoptosis of the cells and leads to antiproliferative effects (Novoa et al. 2001), and EGCG was reported to induce apoptosis of the cells (Okabe et al. 1997; Ahmad et al. 1997; Yang et al. 1998; Okabe et al. 1999). *p21* is an inhibitor of *cyclin-dependent kinase* gene. The upregulation of *p21* gene inhibits proliferation by blocking the cell cycle (El-Deiry et al. 1994). And multiple treatments with EGCG induced enhancement of *GADD153* and *p21* gene expressions (Fig. 2) (Kuzuhara et al. 2007a).

Table 2 Enhanced incorporation of ³H-EGCG by duplicate administrations: the “Fujiki-suganuma effect”

	Total radioactivity (×10 ⁴ dpm)		Fold increase
	Single	Duplicate	
Blood (/ml)	25.13	149.40	×5.9
Brain	3.00	20.29	×6.8
Lung	3.47	16.14	×4.7
Liver	18.94	87.06	×4.6
Kidney	4.68	14.68	×3.1
Spleen	1.33	2.03	×1.5
Pancreas	0.77	3.29	×4.3
Uterus and ovary	2.04	7.23	×3.5
Bladder	0.14	0.74	×5.3
Mammary gland (100 mg)	0.29	0.72	×2.5
Bone (100 mg)	0.96	4.50	×4.7
Skin (100 mg)	3.41	4.91	×1.4





Fig. 2 The induction of *GADD153(L)* and *p21(R)* gene expressions in A549. Multiple treatments with EGCG (filled circle) at 6-h intervals and single treatment with EGCG (open circle)



We think that multiple treatments with EGCG induce synergistic effects on apoptosis in cells and that frequent drinking of green tea plays an important role in cancer prevention.

Inhibitions of in vitro cell growth and TNF- α release, and induction of apoptosis by green tea catechins

Various green tea catechins inhibited the growth of PC-9 cells, dose-dependently, with the order of potency being ECG, EGCG and EGC; EC was not effective (Okabe et al. 1997). Since the activity of EGCG was relatively weak, we compared the IC₅₀ value of EGCG with that of the anticancer drug adriamycin for growth inhibition of two human lung cancer cell lines, PC-9 and PC-14. The IC₅₀ value for adriamycin was 0.17 μ M and that for EGCG was 41.5 μ M (Fig. 3) (Komori et al. 1993a), which means that EGCG is approximately 250-fold less effective than adriamycin.

Although EGCG and green tea extract have very weak anticancer activity, they are unique in having multifunctions: inhibition of cell growth and TNF- α release from the cells induced by okadaic acid, a tumor promoter and

induction of apoptosis with green tea catechins (Table 3) (Suganuma et al. 1999a). EGCG, ECG and EGC were active in all three tests, while EC was inactive. However, cotreatment of EC with EGCG, ECG or EGC synergistically enhanced inhibition of TNF- α release from the cells (Table 4) (Suganuma et al. 1999a). Since we consider TNF- α to be an endogenous tumor promoter, EGCG inhibition of TNF- α release is a key mechanism in cancer prevention (Suganuma et al. 1999b; Fujiki and Suganuma 2011). Synergistic effects were also observed in the inhibition of cell growth and induction of apoptosis. EC obviously increased bioavailability of the catechins, since EC enhanced the incorporation of ³H-EGCG into the cells (Suganuma et al. 1999a). In light of our evidence, we think that usual green tea infusion from the green tea leaves of sencha with warm water is the most appropriate and




Fig. 3 Growth inhibition of two human lung cancer cell lines, PC-9 and PC-14 by EGCG (open circle), green tea extract (filled circle) and adriamycin (times symbol)

Table 3 Three biochemical and biological effects with green tea catechins

Catechins	Inhibition of cell growth (% of control)	Inhibition of TNF- α release IC ₅₀ (μ M)	Induction of apoptosis (A ₄₁₅ nm)
EC (100 ^a , 200 ^b μ M)	97.8 ^b	>500 ^a	0.01 \pm 0.03 ^b
EGCG (100 μ M)	73.3	60	0.52 \pm 0.22
ECG (50 μ M)	62.2	30	0.27 \pm 0.13
EGC (200 μ M)	100.0	ND	0.14 \pm 0.03

ND not determined

^a 100 μ M

^b 200 μ M

Table 4 Synergistic effects by cotreatment of EC with other green tea catechins on inhibition of TNF- α release

	Inhibition of TNF- α release IC ₅₀ (μ M)	
	Without EC	With EC (100 μ M)
		>500
EGCG (100 μ M)	60	15
ECG (50 μ M)	30	7

practical cancer preventive. Precisely how green tea catechins induce multifunctions favorable for cancer prevention is a significant question, and we have presented evidence showing that green tea catechins have potential to act as chemical chaperones (Kuzuhara et al. 2008).

Sealing effects of EGCG and green tea extract

It was exciting for us to find that repeated applications of EGCG inhibited the tumor promotion pathways of both TPA and okadaic acid, although their mechanisms of action are different: TPA and teleocidin bind to the phorbol ester receptor, whereas okadaic acid binds to protein phosphatases 1 and 2A. These two different receptors are present in the membrane and cytosol fractions of mouse skin. When mouse skin was treated with a single application of 5 mg EGCG, both the specific binding of ^3H -TPA and that of ^3H -okadaic acid decreased immediately, within 5–10 min; the levels of the specific binding gradually returned to normal. In the experiments with tumor promotion on mouse skin, EGCG was usually applied 15 min before each application of tumor promoter. Based on the evidence, we think that EGCG treatment inhibited the interaction of tumor promoters with their receptors and that EGCG can interrupt the interaction of ligand with its receptor on cell membrane: This was called the “Sealing Effects of EGCG” (Yoshizawa et al. 1992). Experimentally, EGCG caused reduction in detergent-insoluble membrane domain, that is, a decrease in lipid raft (Fujiki 2005; Adachi et al. 2007), and EGCG also inhibited the activation of specific receptor tyrosine kinases, such as epidermal growth factor receptor, insulin-like growth factor-1 receptor and vascular endothelial growth factor receptor 2 (Shimizu et al. 2011). The results indicate that treatment of cells with EGCG and green tea extract reduced kinase activity, one of the multifunctions of EGCG.

International response to cancer prevention with green tea

In 1991, the International Symposium on Physiological and Pharmacological Effects of *Camellia Sinensis* (Tea) was held at the American Health Foundation in New York. At a press conference, a US scientist said that since animal studies often do not have identical results in humans, more studies were needed before he could recommend green tea for its health effects on humans. However, one of our groups told the press that Japanese drink green tea every day and that green tea could be one of the most practical methods of cancer prevention available. The next day (August 27, 1991), his comments were on the first page of USA Today. We are convinced that his comments in 1991 are now more valid and widely accepted than ever.

Furthermore, in 1994, the New York Times reported the cancer preventive effects of green tea in a story entitled “Green tea: more than just a soothing brew.” This article was based on a report from Shanghai that green tea may reduce the incidence of cancer of the esophagus in humans (June 15, 1994). And in 1995, one of us had the happy opportunity to give a 1-h lecture about our research under the title of “Inhaltsstoffe des grünen Tees, Ein Beispiel für Krebsprevention beim Menschen,” within the framework of Teleakademie von Südwestdeutschen Fernsehen, Baden–Baden in Germany.

Green tea is a popular beverage in Asian countries, and cancer prevention with green tea is now a topic in Western nations. The cover of *Cancer Research* for September 1998 shows the leaves of the green tea plant and its small white flowers, and *Molecular Carcinogenesis* (February, 1999) introduced the results of T. Bowden, in which EGCG in a tea bag inhibited the activation of p38 mitogen-activated protein (MAP) kinase, resulting in the prevention of skin cancer (Fig. 4) (Chen et al. 1999). And in 2011, the 4th World Congress on Tea & Health was held at the Max Delbrück Center for Molecular Medicine in Berlin Buch, introducing new challenging applications of EGCG for neurodegenerative diseases.

Prospective cohort study in Saitama Prefecture

The cancer preventive effect of green tea was first confirmed through a prospective cohort study with humans in Saitama Prefecture. In 1986, we first surveyed 8,552 individuals aged over 40 on their living habits, including their daily consumption of green tea. During the 10 years after 1986, a total of 419 cancer patients, 244 males and 175 females, were detected (Nakachi et al. 2000). Female and male cancer patients were divided into three groups based on consumption of green tea per day: under 3 cups, 4–9




Fig. 4 The covers of *Cancer Research* (L) and *Molecular Carcinogenesis* (R)

Table 5 Average age at cancer onset and daily green tea consumption

Gender	Average age at cancer onset and consumption of green tea per day (cups)		
	≤3	4–9	≥10
Female (175)	67.0 ± 1.7 (28.0 %)	66.4 ± 1.3 (58.3 %)	74.3 ± 2.2 (13.7 %)
Male (244)	65.0 ± 1.5 (24.2 %)	67.2 ± 1.0 (46.7 %)	68.2 ± 1.1 (29.1 %)



cups and over 10 cups (Table 5). Next, the average age at cancer onset of all these cancer patients was obtained from clinical documents at hospitals. The average age at cancer onset and daily green tea consumption per patient were studied.

The cancer onset in female patients who had consumed over 10 Japanese-size cups (120 ml/cup) of green tea per day was 7.3 years later than that of female patients who had consumed less than three cups per day. The cancer onset in male patients who had consumed over 10 cups of green tea per day was 3.2 years later than that of patients who had consumed less than three cups per day (Table 5) (Nakachi et al. 2000). The difference between females and males is partly due to higher tobacco consumption by males. It is important to note that the delay of cancer onset is significant evidence of primary cancer prevention in humans. Key factors in primary cancer prevention include: tamoxifen for breast cancer (Fisher et al. 1998), cessation of smoking, diet modifications and physical activity. In addition, the cohort study first demonstrated that consumption of 10 cups of green tea per day is an additional significant factor in primary cancer prevention for the general population. Since the consumers of over 10 cups of green tea per day were only 13.7 % for female patients and 29.1 % for male patients, it would be possible to increase consumption to over 10 cups per day with green tea tablets, resulting in effective primary cancer prevention (Fujiki et al. 2002).

We next studied the reduced relative risk in human organs by high consumption of green tea in Saitama Prefecture. Green tea most significantly prevented lung cancer—a relative risk of 0.33 with over 10 cups of green tea per day. High consumption of green tea also prevented cancers of the colorectum, liver and stomach, in that order (Fig. 5) (Nakachi et al. 2000). It is important to note that consumption of less than ten cups is not effective as a preventive tool, suggesting that there is a minimum effective amount of green tea catechins for cancer prevention. In addition to the cancer preventive effects, the prospective cohort study provided other significant results. Increased green tea consumption was associated with decreased serum total cholesterol, decreased triglyceride levels and decreased atherogenic index. Moreover, the prevalence rates of cardiovascular disease and diabetes mellitus were significantly lower among the population consuming over 10 cups of green tea per day (Imai et al. 1997). Since green tea apparently prevents many major

lifestyle-related diseases, including cancer, cardiovascular disease and diabetes mellitus, we reported that among women, the mean age at death for those consuming over 10 cups per day was 6 years later than that for those consuming less than 3 cups, suggesting that green tea reduces the risk of lifestyle-related diseases, resulting in a longer life span (Imai et al. 1997; Nakachi et al. 2000; Sueoka et al. 2001). All the results of this study indicate that the more green tea we drink, the higher cancer preventive activity we get in the target organs. Of course, cancer prevention for general population before cancer onset is a matter of individual responsibility, not the duty of clinicians. We call this the first stage of cancer prevention with green tea (Fig. 6).

Our cohort study was conducted in Saitama Prefecture, one of the main tea-producing areas in Japan. Other investigators have reported non-preventive effects of green tea against human cancers, such as no association between the consumption of green tea and the risk of gastric cancer

**Fig. 5** Reduced relative risk of cancer in human organs with a high consumption of green tea. ▨ ≤3, ▤ 4–9, ▥ ≥10**Fig. 6** Two stages of cancer prevention with green tea

in Miyagi Prefecture (Tsubono et al. 2001), and no association between green tea drinking and the risk of breast cancer in Japan (Iwasaki et al. 2010). It is important to note that various kinds of green tea are produced in Japan, such as sencha, hojicha and oolong tea. Sencha contains 13.6 % green tea catechins, hojicha made from roasted lowest grade of tea leaf contains only 2.9 %, and oolong tea 7.6 %. This clearly shows that the results of epidemiological study are dependent on the kinds of green tea consumed and the quality of green tea catechins. Thus, an ingested amount of green tea catechins is a limiting factor for demonstrating the preventive or non-preventive effects in a cohort study.

In addition, an interesting study reported that no association between green tea consumption and breast cancer incidence was found in the cohort studies of the United States, but that increased green tea consumption (more than three cups a day) was inversely associated with breast cancer recurrence (Pooled RR = 0.73, 95 % CI: 0.56-0.96) (Ogunleye et al. 2010). In 1998, we previously reported the decreased recurrence of human breast cancer with increased consumption of green tea, on the basis of results obtained from 472 cancer patients. Stages I and II cancer patients consuming over five cups of green tea per day (average 8 cups) showed a lower recurrence rate, 16.7 %, and a longer disease-free period, 3.6 years, than those consuming less than four cups (average 2 cups) per day, 24.3 % and 2.8 years (Nakachi et al. 1998). The results showed that drinking an average 8 cups of green tea per day prior to cancer onset results in more hopeful prognosis for breast cancer patients.

Preclinical safety trials for 10 cups of green tea

Next, we will think about cancer prevention for patients following cancer treatment, since surviving cancer patients are seriously looking for preventives. Our epidemiological results showed that the effective cancer preventive amount is also 10 Japanese-size cups of green tea per day, corresponding to 2.5 g green tea extract (Nakachi et al. 2000). To facilitate consumption, tablets of green tea extract are produced by the Tea Institute of Saitama Prefecture. A tablet dissolved in warm water becomes usual Saitama tea (Fujiki et al. 2001). 10 cups of green tea per day may seem to be a large amount, but the results of Phase I clinical trials with green tea extract conducted in USA reported that the maximum tolerated dose corresponds to 21–24 Japanese-size cups per day (Pisters et al. 2001), or even higher amounts (Laurie et al. 2005). Therefore, we think 10 cups of green tea is tolerable for most people.

Before going on to a clinical trial, we first asked some 102 healthy and voluntary citizens to consume this amount for 3 months, with informed consent. Group A of 51




Fig. 7 Preclinical safety trials among 102 healthy and voluntary citizens of Saitama Prefecture for 6 months. 10 Cups of daily green tea beverage fortified by green tea tablets for 3 months

volunteers took 10 cups of daily green tea beverage supplemented with green tea tablets for the first 3 months, and group B took the same amount for the next 3 months. During the 6-month trial, all of the 102 volunteers answered questionnaires and their blood samples were biochemically examined three times (Fig. 7) (Fujiki et al. 2001). The blood examination did not show any serious side effects, so important answers to just two of the questions are presented. To the question, “Did you experience any discomfort from taking green tea tablets?”, 93 % of the participants said that they were able to continue drinking green tea and taking green tea tablets. For another question, “Did you observe any changes in living habits or meals?”, Over 90 % of the participants found their living habits or appetite unaffected by 10 cups of green tea daily.

However, about 50 % of the volunteers experienced very mild temporary disorders due to the caffeine in green tea extract. It is reported that the maximum tolerable dose of caffeine is about 1.0 g/day, and even though 10 cups do not contain total 1.0 g caffeine, the Tea Institute developed a new method to reduce the caffeine by washing the green tea leaves with warm water. They reduced the caffeine content from 5 % to less than 3 %, without using an organic solution. It is important to note that the tablets are a dried green tea infusion, not a powder of green tea leaves. The tablets are not controlled by the Pharmaceutical Affairs Act in Japan because they belong to the category of green tea beverage, not a drug.

Significant factors for studying cancer prevention with humans

Before going on to human studies, we had to clarify the following factors. (1) It is necessary to have evidence that green tea is non-toxic for humans and that there is bio-availability in rodents (Suganuma et al. 1998). (2) We obtained the minimum effective amounts of green tea per day for humans from the results of human epidemiological study (Nakachi et al. 2000). (3) Collaboration of healthy volunteers with informed consent was established (Fujiki et al. 2001). (4) Clinicians developed an established clinical biomarker for target organ (Shimizu et al. 2008).

(5) Green tea tablets should be quality controlled, and the absence of pesticides and herbicides confirmed (Fujiki et al. 2001). (6) We were fortunate to be able to collaborate with the Tea Institute of Saitama Prefecture, which provided green tea extract and tablets. (7) Results should be published in reviewed international journals (Nakachi et al. 2000; Shimizu et al. 2008).

Double-blind randomized clinical phase II prevention trial for recurrence of colorectal adenomas

Table 6 shows the protocol for phase II prevention trial of colorectal adenoma recurrence with green tea extract, with informed consent, which was conducted at Gifu University, Department of Medicine and its related hospitals (Shimizu et al. 2008). One green tea tablet (GTE, 500 mg) contains 52.5 mg EGCG, 34.6 mg EGC, 11.1 mg ECG, 12.3 mg EC and 15.7 mg caffeine, approximately equivalent to two Japanese-size cups of green tea. Colorectal adenomas were removed by endoscopic polypectomy at the first colonoscopy, and 12 months later, the absence of polyps was confirmed by second colonoscopy. The patients were then double-blind randomized into two groups: one group maintained their usual daily consumption of green tea beverage without a placebo, and the other group (GTE group) took 10 cups of green tea, supplemented with six green tea tablets, daily for 12 months. The incidence of recurrent adenomas was determined by end-point colonoscopy 12 months later. The recurrence rate of the control group was 31 % and that of the GTE group was 15 % (Shimizu et al. 2008). The chi-square test, p was <0.05 (Table 6). In addition, the size of recurrent adenomas was smaller in the GTE group than in the control group. Thus, drinking 10 Japanese-size cups of green tea, supplemented with green tea tablets, 50 %, significantly prevented

recurrence of colorectal adenomas in patients (Shimizu et al. 2008).


It is important to note that we did not use placebo in our phase II prevention trial because green tea is a daily beverage of Japanese, so, we aimed to raise the consumption to a target level, with supplemental tablets of green tea extract. The patients in the trial were told to take a minimum effective amount of green tea catechins. A randomized, placebo-controlled multicentre trial was recently begun to investigate the effects of diet supplementation with green tea extract containing 300 mg EGCG on the recurrence of colon adenomas for patients who have undergone polypectomy for colon polyps in Germany (Stingl et al. 2011).

Moreover, the prevention of prostate cancer development in patients with high-grade prostate intraepithelial neoplasia (PIN) using capsules of green tea catechins was confirmed in Italy. In the group using green tea capsules, only one tumor was diagnosed among 30 patients, whereas nine cancers were found among 30 placebo-controlled patients. Green tea catechins reduced incidence of prostate cancer from 30 to 3 % (Bettuzzi et al. 2006). In the United States, a phase I trial of green tea extract was first conducted at two Institutions (Pisters et al. 2001), and a phase II randomized, placebo-controlled trial of green tea extract in patients with high-risk oral premalignant lesions revealed that the two higher doses (750 and 1,000 mg/m²) of green tea extract produced a trend toward a greater clinical response in association with a high baseline level and downregulation of angiogenic stromal vascular endothelial growth factor (Tsao et al. 2009).

Necessity of cancer prevention in our daily lives

Tumor promotion in human cancer is now understood as a disease affected by TNF- α upregulation and NF- κ B

Table 6 Phase II prevention trial of colorectal adenoma recurrence with green tea extract in patients, with no polyps after polypectomy



Control group 65	Daily consumption of green tea without placebo
GTE group 60	Daily 10 Cups of green Tea + GTE

Groups (cases)	Recurrence rate (%)	Average no. of polyps/patient
Control (20/65)	31.0	0.43
GTE (9/60)	15.0 ($p < 0.05$)	0.20

activation (Komori et al. 1993b; Suganuma et al. 1999b; Ben-Neriah and Karin 2011), induced by pro-inflammatory cytokines such as TNF- α , IL-1 and IL-6 (Suganuma et al. 2002). Pro-inflammatory cytokines are easily induced in humans by bacterial focal infections (Fujiki et al. 2004; Kuzuhara et al. 2007b); Cytokines and chemokines are numerous in aged people; and so, humans are always at risk of tumor promotion. With these physiological conditions, we need to establish our own cancer prevention strategy, based on reduction in TNF- α and inactivation of NF- κ B. For this purpose, we believe cancer prevention with green tea is the most appropriate preventive, since it is effective on a wide range of target organs and has no toxic effects. Green tea can be ingested by everybody in the world.

Due to advancements in diagnosis and treatment, there are many healthy surviving cancer patients in Japan. They are seriously looking for drugs and preventives that will prevent recurrence, second primary tumors and metastasis. For these patients, green tea supplemented with green tea tablets is vital following cancer treatment. We call this the two stages of cancer prevention with green tea (Fig. 6) (Fujiki 1999; Fujiki et al. 2002). The left part of Fig. 6, the first stage, deals with primary cancer prevention, and the right part of Fig. 6, the second stage, shows tertiary cancer prevention.

New cancer treatment strategy using the combination of green tea catechins and anticancer drugs

Considering the cancer prevention for patients following cancer treatment (Fig. 6), anticancer drugs may be administered to Japanese cancer patients who consume green tea every day. So we raised the following questions: Is it advantageous for Japanese cancer patients to take cancer preventive green tea catechins and anticancer drugs together? Does the combination have the potential to enhance efficacy and decrease adverse effects of anticancer drugs? What kinds of anticancer drugs and compounds can work with green tea catechins? The names of the compounds are presented: gefitinib, 5-fluorouracil, taxol, doxorubicin, erlotinib, antiestrogen, COX inhibitors, sulindac and celecoxib, retinoids and curcumin.

We had two significant results. The combination of green tea extract with sulindac enhanced the inhibition of tumor development in multiple intestinal neoplasia (Min) mice (Suganuma et al. 2001). Min mice have a germline mutation of *APC* gene and develop intestinal tumors similar to those of familiar adenomatous polyposis, FAP patients. Intestinal tumors are seen as blue spots on the organ by staining with methylen blue. The combination reduced the number of tumors per mouse from 72.3 to 32.0, a decrease of 55.7 %, $p < 0.05$ (Suganuma et al. 2001). The second result was as follows: the combination of green

tea extract with celecoxib enhanced the inhibition of tumor development in the lungs of A/J mice induced by 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK), a carcinogen. The combination reduced the number of tumor-bearing mice from 100 to 73.3 %, a decrease of 26.7 %, $p < 0.05$ (Suganuma et al. 2011).

To understand the molecular mechanisms involved in synergistic enhancement by the combination of EGCG with sulindac, we first studied the upregulation of two gene expressions, *GADD153* and *p21* genes, and as previously reported, their inductions of gene expression were enhanced to be about 12-fold and 3-fold, respectively, (Okabe et al. 2001; Fujiki and Suganuma 2002). These gene expressions were not affected by treatment with either EGCG alone or sulindac alone. The combination of EGCG with sulindac induced apoptosis in PC-9 cells 10.5-fold stronger than EGCG alone or sulindac alone, and the combination of EGCG with celecoxib induced apoptosis in PC-9 cells 14.9-fold stronger than EGCG alone or celecoxib alone (Table 7) (Suganuma et al. 2006). The results show that the combination additionally induces new gene expressions that are not induced by EGCG alone or COX inhibitors alone. A schematic illustration of the combination is shown in Fig. 8. The combinations indicate the presence of a new therapeutic activity that will greatly expand cancer treatment and also bring successful tertiary cancer prevention for patients with good prognosis (Suganuma et al. 2011).

Our definition of cancer prevention with green tea

The administration of cancer preventives delays the carcinogenic processes in humans, no matter when the carcinogenesis starts, thereby blocking the appearance of clinical symptoms (Fujiki et al. 2002).



Afterword

Strong support by Professor James Watson

In May 2002, the International Conference on Green Tea and Cancer, A Critical Review, was held at the Banbury Center, Cold Spring Harbor Laboratory in New York. This Conference was strongly supported by the Director, J. Watson. After the Conference at the Banbury Center, Dr. Watson listened closely while one of the organizers noted the success of the Conference. The organizer told him that green tea clearly prevented cancer in rodents, whereas human epidemiological studies had proved inconclusive. Dr. Watson immediately answered that since green tea prevented carcinogenesis in rodents, it should prevent it in humans as well, since humans and mice have

Table 7 Synergistic induction of apoptosis by combination of EGCG with COX inhibitors

	Induction of apoptosis (% of apoptotic cells)	
	Without	With EGCG
Human lung cancer cell line PC-9		
Control	2.0	6.3
Sulindac	4.1	42.9 (×10.5)
Celecoxib	3.0	44.6 (×14.9)

**Fig. 8** Combination of green tea with COX inhibitor

similar genomes. It was wonderful encouragement for green tea scientists like us, because we got the impression that a world-famous molecular biologist now believes in the cancer preventive effects of green tea for humans. Seven years later, we successfully confirmed the prevention of colon adenomas in humans by green tea supplemented with green tea tablets.

The history of green tea in Japan

Drinking green tea is really part of Japanese culture. The Japanese Zen priest Eisai returned to Japan after 4 years training as Zen priest in China, carrying some seeds of green tea, as a medicine. When the third Shogun became ill, Eisai wrote a book entitled “Kitusa yohjohki” (Maintaining Health by Drinking Green Tea), which was

presented to the Shogun together with green tea in the year 1211. The Shogun’s illness, probably a disorder of the digestive tract, was cured soon, and this is the first case of therapeutic effects with green tea in Japan.

Numerous illnesses of course existed 800 years ago, but cancer and aging-associated diseases did not, because the average life span was only 40 years at that time. Thus, our study for the first time demonstrated that green tea prevents a lifestyle-related disease. Eisai concluded his book this way: I hope an excellent doctor will elucidate the mechanism of green tea action in the near future. We believe our research project with green tea is a present from Eisai.

Acknowledgments To Prof. Erich Hecker, we express gratitude for his fruitful discussion. We thank Prof. Takuo Okuda for providing numerous polyphenols including (–)-epigallocatechin gallate (EGCG) and penta-*O*-galloyl-β-*D*-glucose (5GG), Prof. Takashi Sugimura for his encouragement during the course of this work, and our Japanese colleagues cited in our references for their research. This work was supported by Japan Society for the Promotion of Science: Scientific Research on Priority Areas for Cancer Research from the Ministry of Education, Culture, Sports, Science and Technology, Japan; Comprehensive Research on Aging and Health; and Cancer Research from the Ministry of Health, Labor, and Welfare, Japan; Selective Applied and Developed Research, and Green Tea Extracts Research Development for Cancer Prevention by the Department of Agriculture and Forests and the Department of Health and Human Services of Saitama Prefecture, Japan; and the Smoking Research Fund.

Conflict of interest We declare that we have no conflict of interest.

References

- Adachi S, Nagao T, Ingolfsson HI, Maxfield FR, Andersen OS, Kopelovich L, Weinstein IB (2007) The inhibitory effect of (–)-epigallocatechin gallate on activation of the epidermal growth factor receptor is associated with altered lipid order in HT29 colon cancer cells. *Cancer Res* 67:6493–6501
- Ahmad N, Feyes DK, Nieminen AL, Agarwal R, Mukhtar H (1997) Green tea constituent epigallocatechin-3-gallate and induction of apoptosis and cell cycle arrest in human carcinoma cells. *J Natl Cancer Inst* 89:1881–1886
- Ben-Neriah Y, Karin M (2011) Inflammation meets cancer, with NF- κ B as the matchmaker. *Nat Immunol* 12:715–723
- Bettuzzi S, Brausi M, Rizzi F, Castagnetti G, Peracchia G, Corti A (2006) Chemoprevention of human prostate cancer by oral administration of green tea catechins in volunteers with high-grade prostate intraepithelial neoplasia: a preliminary report from a one-year proof-of-principle study. *Cancer Res* 66:1234–1240
- Chen W, Dong Z, Valcic S, Timmermann BN, Bowden GT (1999) Inhibition of ultraviolet B-induced *c-fos* gene expression and p38 mitogen-activated protein kinase activation by (–)-epigallocatechin gallate in a human keratinocyte cell line. *Mol Carcinog* 24:79–84
- Conney AH, Wang ZY, Huang MT, Ho CT, Yang CS (1992) Inhibitory effect of oral administration of green tea on tumorigenesis by ultraviolet light, 12-*O*-tetra decanoylphorbol-13-acetate and *N*-nitrosodiethylamine in mice. CRC Press, Boca Raton, pp 361–373
- El-Deiry WS, Harper JW, O'Connor PM, Velculescu VE, Canman CE, Jackman J, Pietenpol JA, Burrell M, Hill DE, Wang Y, Wiman KG, Mercer WE, Kastan MB, Kohn KW, Elledge SJ, Kinzler KW, Vogelstein B (1994) *WAF1/CIP1* is induced in *p53*-mediated G₁ arrest and apoptosis. *Cancer Res* 54:1169–1174
- Fisher B, Costantino JP, Wickerham DL, Redmond CK, Kavanah M, Cronin WM, Vogel V, Robidoux A, Dimitrov N, Atkins J, Daly M, Wieand S, Tan-Chiu E, Ford L, Wolmark N, other National Surgical Adjuvant Breast and Bowel Project Investigators (1998) Tamoxifen for prevention of breast cancer. Report of the national surgical adjuvant breast and bowel project P-1 study. *J Natl Cancer Inst* 90:1371–1388
- Fujiki H (1999) Two stages of cancer prevention with green tea. *J Cancer Res Clin Oncol* 125:589–597
- Fujiki H (2005) Green tea: health benefits as cancer preventive for humans. *Chem Rec* 5:119–132
- Fujiki H, Okuda T (1992) (–)-Epigallocatechin gallate. *Drugs Future* 17:462–464
- Fujiki H, Suganuma M (2002) Green tea and cancer prevention. *Proc Jpn Acad* 78(B):263–270
- Fujiki H, Suganuma M (2011) Tumor promoters-microcystin-LR, nodularin and TNF- α and human cancer development. *Anticancer Agents Med Chem* 11:4–18
- Fujiki H, Suganuma M, Okabe S, Komori A, Sueoka E, Sueoka N, Kozu T, Sakai Y (1996) Japanese green tea as a cancer preventive in humans. *Nutr Rev* 54:S67–S70
- Fujiki H, Suganuma M, Okabe S, Sueoka E, Sueoka N, Fujimoto N, Goto Y, Matsuyama S, Imai K, Nakachi K (2001) Cancer prevention with green tea and monitoring by a new biomarker, hnRNP B1. *Mutat Res* 480–481:299–304
- Fujiki H, Suganuma M, Imai K, Nakachi K (2002) Green tea: cancer preventive beverage and/or drug. *Cancer Lett* 188:9–13
- Fujiki H, Takeuchi H, Nishitani N, Yamanaka H, Suzuki K, Kurusu M, Suganuma M (2004) Carcinogenic potential of tobacco tar-resistant *Staphylococcus aureus* in buccal cavity. *J Cancer Res Clin Oncol* 130:301–305
- Fujita Y, Yamane T, Tanaka M, Kuwata K, Okuzumi J, Takahashi T, Fujiki H, Okuda T (1989) Inhibitory effect of (–)-epigallocatechin gallate on carcinogenesis with *N*-ethyl-*N'*-nitro-*N*-nitrosoguanidine in mouse duodenum. *Jpn J Cancer Res* 80:503–505
- Gupta S, Hastak K, Ahmad N, Lewin JS, Mukhtar H (2001) Inhibition of prostate carcinogenesis in TRAMP mice by oral infusion of green tea polyphenols. *Proc Natl Acad Sci USA* 98:10350–10355
- Hecker E, Adolf W, Hergenbahn R, Schmidt R, Sorg B (1984) Irritant diterpene ester promoters of mouse skin: contributions to etiologies of environmental cancer and to biochemical mechanisms of carcinogenesis. In: Fujiki H, Hecker E, Moore RE, Sugimura T, Weinstein IB (eds) Cellular interactions by environmental tumor promoters. Jpn Sci Soc Press/VNU Science Press, BV, Tokyo/Utrecht, pp 3–36
- Imai K, Suga K, Nakachi K (1997) Cancer-preventive effects of drinking green tea among a Japanese population. *Prev Med* 26:769–775
- Iwasaki M, Inoue M, Sasazuki S, Sawada N, Yamaji T, Shimazu T, Willett WC, Tsugane S (2010) Green tea drinking and subsequent risk of breast cancer in a population to based cohort of Japanese women. *Breast Cancer Res* 12:R88. (Epub ahead of print)
- Komori A, Yatsunami J, Okabe S, Abe S, Hara K, Suganuma M, Kim SJ, Fujiki H (1993a) Anticarcinogenic activity of green tea polyphenols. *Jpn J Clin Oncol* 23:186–190
- Komori A, Yatsunami J, Suganuma M, Okabe S, Abe S, Sakai A, Sasaki K, Fujiki H (1993b) Tumor necrosis factor acts as a tumor promoter in BALB/3T3 cell transformation. *Cancer Res* 53:1982–1985
- Kuzuhara T, Sei Y, Yamaguchi K, Suganuma M, Fujiki H (2006) DNA and RNA as new binding targets of green tea catechins. *J Biol Chem* 281:17446–17456
- Kuzuhara T, Tanabe A, Sei Y, Yamaguchi K, Suganuma M, Fujiki H (2007a) Synergistic effects of multiple treatments, and both DNA and RNA direct bindings on, green tea catechins. *Mol Carcinog* 46:640–645
- Kuzuhara T, Suganuma M, Kurusu M, Fujiki H (2007b) *Helicobacter pylori*-secreting protein Tipz is a potent inducer of chemokine gene expressions in stomach cancer cells. *J Cancer Res Clin Oncol* 133:287–296
- Kuzuhara T, Suganuma M, Fujiki H (2008) Green tea catechin as a chemical chaperone in cancer prevention. *Cancer Lett* 261:12–20
- Laurie SA, Miller VA, Grant SC, Kris MG, Ng KK (2005) Phase I study of green tea extract in patients with advanced lung cancer. *Cancer Chemother Pharmacol* 55:33–38
- Nakachi K, Suemasu K, Suga K, Takeo T, Imai K, Higashi Y (1998) Influence of drinking green tea on breast cancer malignancy among Japanese patients. *Jpn J Cancer Res* 89:254–261
- Nakachi K, Matsuyama S, Miyake S, Suganuma M, Imai K (2000) Preventive effects of drinking green tea on cancer and cardiovascular disease: epidemiological evidence for multiple targeting prevention. *BioFactors* 13:49–54
- Narisawa T, Fukaura Y (1993) A very low dose of green tea polyphenols in drinking water prevents *N*-methyl-*N*-nitrosourea-induced colon carcinogenesis in F344 rats. *Jpn J Cancer Res* 84:1007–1009
- NCI, Dcpc, Chemoprevention Branch and Agent Development Committee (1996) Clinical development plan:tea extracts green tea polyphenols epigallocatechin gallate. *J Cell Biochem* 26S:236–257
- Nishida H, Omori M, Fukutomi Y, Ninomiya M, Nishiwaki S, Suganuma M, Moriaki H, Muto Y (1994) Inhibitory effects of (–)-epigallocatechin gallate on spontaneous hepatoma in C3H/HeNcrj mice and human hepatoma-derived PCL/PRF/5 cells. *Jpn J Cancer Res* 85:221–225

- Novoa I, Zeng H, Harding HP, Ron D (2001) Feedback inhibition of the unfolded protein response by *GADD34*-mediated dephosphorylation of eIF2 α . *J Cell Biol* 153:1011–1022
- Ogunleye AA, Xue F, Michels KB (2010) Green tea consumption and breast cancer risk or recurrence: a meta-analysis. *Breast Cancer Res Treat* 119:477–484
- Okabe S, Suganuma M, Hayashi M, Sueoka E, Komori A, Fujiki H (1997) Mechanisms of growth inhibition of human lung cancer cell line, PC-9, by tea polyphenols. *Jpn J Cancer Res* 88:639–643
- Okabe S, Ochiai Y, Aida M, Park K, Kim SJ, Nomura T, Suganuma M, Fujiki H (1999) Mechanistic aspects of green tea as a cancer preventive: effect of components on human stomach cancer cell lines. *Jpn J Cancer Res* 90:733–739
- Okabe S, Fujimoto N, Sueoka N, Suganuma M, Fujiki H (2001) Modulation of gene expression by (–)-epigallocatechin gallate in PC-9 cells using a cDNA expression array. *Biol Pharm Bull (Tokyo)* 24:883–886
- Okuda T, Mori K, Hayatsu H (1984) Inhibitory effect of tannins on direct-acting mutagens. *Chem Pharm Bull (Tokyo)* 32:3755–3758
- Okuda T, Mori K, Hatano T (1985) Relationship of the structure of tannins to the binding activities with hemoglobin and methylene blue. *Chem Pharm Bull (Tokyo)* 33:1424–1433
- Pisters KM, Newman RA, Coldman B, Shin DM, Khuri FR, Hong WK, Glisson BS, Lee JS (2001) Phase I trial of oral green tea extract in adult patients with solid tumors. *J Clin Oncol* 19:1830–1838
- Shimizu M, Fukutomi Y, Ninomiya M, Nagura K, Kato T, Araki H, Suganuma M, Fujiki H, Moriwaki H (2008) Green tea extracts for the prevention of metachronous colorectal adenomas: a pilot study. *Cancer Epidemiol Biomarkers Prev* 17:3020–3025
- Shimizu M, Adachi S, Masuda M, Kozawa O, Moriwaki H (2011) Cancer chemoprevention with green tea catechins by targeting receptor tyrosine kinases. *Mol Nutr Food Res* 55:832–843
- Sporn MB, Dunlop NM, Newton DL, Smith JM (1976) Prevention of chemical carcinogenesis by vitamin A and its synthetic analogs (retinoids). *Fed Proc* 35:1332–1338
- Stingl JC, Ettrich T, Muche R, Wiedom M, Brockmüller J, Seeringer A, Seufferlein T (2011) Protocol for minimizing the risk of metachronous adenomas of the colorectum with green tea extract (MIRACLE): a randomized controlled trial of green tea extract versus placebo for nutripredvention of metachronous colon adenomas in the elderly population. *BMC Cancer* 11:360
- Sueoka N, Suganuma M, Sueoka E, Okabe S, Matsuyama S, Imai K, Nakachi K, Fujiki H (2001) A new function of green tea: prevention of lifestyle-related diseases. *Ann N Y Acad Sci* 928:274–280
- Suganuma M, Fujiki H, Suguri H, Yoshizawa S, Hirota M, Nakayasu M, Ojika M, Wakamatsu K, Yamada K, Sugimura T (1988) Okadaic acid: an additional non-phorbol-12-tetradecanoate-13-acetate-type tumor promoter. *Proc Natl Acad Sci USA* 85:1768–1771
- Suganuma M, Okabe S, Oniyama M, Tada Y, Ito H, Fujiki H (1998) Wide distribution of [³H](–)-epigallocatechin gallate, a cancer preventive tea polyphenol, in mouse tissue. *Carcinogenesis* 19:1771–1776
- Suganuma M, Okabe S, Kai Y, Sueoka N, Sueoka E, Fujiki H (1999a) Synergistic effects of (–)-epigallocatechin gallate with (–)-epicatechin, sulindac, or tamoxifen on cancer-preventive activity in the human lung cancer cell line PC-9. *Cancer Res* 59:44–47
- Suganuma M, Okabe S, Marino MW, Sakai A, Sueoka E, Fujiki H (1999b) Essential role of tumor necrosis factor α (TNF- α) in tumor promotion as revealed by TNF- α -deficient mice. *Cancer Res* 59:4516–4518
- Suganuma M, Ohkura Y, Okabe S, Fujiki H (2001) Combination cancer chemoprevention with green tea extract and sulindac shown in intestinal tumor formation in Min mice. *J Cancer Res Clin Oncol* 127:69–72
- Suganuma M, Okabe S, Kurusu M, Iida N, Ohshima S, Saeki Y, Kishimoto T, Fujiki H (2002) Discrete roles of cytokines, TNF- α , IL-1, IL-6 in tumor promotion and cell transformation. *Int J Oncol* 20:131–136
- Suganuma M, Kurusu M, Suzuki K, Tasaki E, Fujiki H (2006) Green tea polyphenol stimulates cancer preventive effects of celecoxib in human lung cancer cells by upregulation of *GADD153* gene. *Int J Cancer* 119:33–40
- Suganuma M, Saha A, Fujiki H (2011) New cancer treatment strategy using combination of green tea catechins and anticancer drugs. *Cancer Sci* 102:317–323
- Taniguchi S, Fujiki H, Kobayashi H, Go H, Miyado K, Sadano H, Shimokawa R (1992) Effect of (–)-epigallocatechin gallate, the main constituent of green tea, on lung metastasis with mouse B16 melanoma cell lines. *Cancer Lett* 65:51–54
- Tsao AS, Liu D, Martin J, Tang X-m, Lee JJ, El-Naggar AK, Wistuba I, Culotta KS, Mao L, Gillenwater A, Sagesaka YM, Hong WK, Papadimitrakopoulou V (2009) Phase II randomized, placebo-controlled trial of green tea extract in patients with high-risk oral premalignant lesions. *Cancer Rev Res* 2:931–941
- Tsubono Y, Nishino Y, Komatsu S, Hsieh C-C, Kanemura S, Tsuji I, Nakatsuka H, Fukao A, Satoh H, Hisamichi S (2001) Green tea and the risk of gastric cancer in Japan. *N Engl J Med* 344:632–636
- Vogelstein B, Fearon ER, Hamilton SR, Kern SE, Preisinger AC, Leppert M, Nakamura Y, White R, Smith AMM, Bos JL (1988) Genetic alterations during colorectal-tumor development. *N Engl J Med* 319:525–532
- Wang ZY, Hong JY, Huang MT, Reuhl K, Conney AH, Yang CS (1992) Inhibition of *N*-nitrosodiethylamine- and 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone-induced tumorigenesis in A/J mice by green tea and black tea. *Cancer Res* 52:1943–1947
- Yamane T, Hagiwara N, Tateishi M, Akachi S, Kim M, Okuzumi J, Kitao Y, Inagake M, Kuwata K, Takahashi T (1991) Inhibition of azoxymethane-induced colon carcinogenesis in rat by green tea polyphenol fraction. *Jpn J Cancer Res* 82:1336–1339
- Yamane T, Takahashi T, Kuwata K, Oya K, Inagake M, Kitao Y, Suganuma M, Fujiki H (1995) Inhibition of *N*-methyl-*N'*-nitro-*N*-nitrosoguanidine-induced carcinogenesis by (–)-epigallocatechin gallate in the rat glandular stomach. *Cancer Res* 55:2081–2084
- Yang CS, Wang ZY (1993) Tea and cancer. *J Natl Cancer Inst* 85:1038–1049
- Yang GY, Liao J, Kim K, Yurkow EJ, Yang CS (1998) Inhibition of growth and induction of apoptosis in human cancer cell lines by tea polyphenols. *Carcinogenesis* 19:611–616
- Yoshizawa S, Horiuchi T, Fujiki H, Yoshida T, Okuda T, Sugimura T (1987) Antitumor promoting activity of (–)-epigallocatechin gallate, the main constituent of “tannin” in green tea. *Phytother Res* 1:44–47
- Yoshizawa S, Horiuchi T, Suganuma M, Nishiwaki S, Yatsunami J, Okabe S, Okuda T, Muto Y, Frenkel K, Troll W, Fujiki H (1992) Penta-*O*-galloyl- β -D-glucose and (–)-epigallocatechin gallate. *Am Chem Soc*:316–325