

Metabolic Adaptations to Weight Loss

Kevin D. Hall 

Does losing weight slow down metabolism? Absolutely. Does metabolism decrease more than expected? It depends on how you set your expectations.

Larger people typically burn more calories; therefore, energy expenditure (EE) declines as people lose weight. Quantifying the expected decline in EE with weight loss requires a mathematical model relating EE to changes in body composition. Deviations between the expected and observed decreases in EE define the degree of metabolic adaptation.

Linear regression is often used to model how resting metabolic rate (RMR) relates to fat-free mass (FFM) and fat mass (FM) in people maintaining their habitual weight. The resulting equations vary depending on the cohort, measurement methods, and the variables included in the model. As an indication of the validity of the regression model, the best-fit coefficients for FFM and FM should be similar to the values calculated by a physiologically based model that accounts for how different organs, varying in size and metabolic rate (1), are related to FFM (19 kcal/kg/d) and FM (4.5 kcal/kg/d) (2). Furthermore, the regression model should explain much of the EE variance between people prior to weight loss because large residuals limit the power to detect metabolic adaptation.

In the current issue of *Obesity*, Wolfe et al. (3) addressed the question of metabolic adaptation by developing regression equations relating RMR and total EE to FFM in patients prior to various bariatric surgeries to predict RMR and total EE after 6 and 24 months. Metabolic adaptation was present at 6 months while patients were still losing weight, but RMR and total EE normalized 2 years after the surgery after accounting for the observed FFM changes.

These results support a growing consensus that metabolic adaptation occurs during periods of negative energy balance. However, much debate surrounds whether metabolic adaptation persists after body weight stabilizes at a lower level. A previous study was similar to Wolfe et al. (3) in that a significant metabolic adaptation 6 months after Roux-en-Y gastric bypass (RYGB) dissipated after 1 year (4). Interestingly, similar weight losses induced by 7 months of caloric restriction and exercise during “The Biggest Loser” competition resulted in comparable metabolic adaptation as 6 months after RYGB (4). However, unlike the matched RYGB group, metabolic adaptation persisted 6 years after “The Biggest Loser” competition despite contestants regaining two-thirds of the lost weight (5), possibly because of increases in physical activity (6).

Previous studies have found persistent metabolic adaptation 1 year or more after bariatric surgery (7-11) or diet-induced weight loss (12,13) after body weight had stabilized. Others failed to detect significant metabolic adaptation after achieving a stable lower weight (14-16), and a large recent study suggested that RMR adjusted for body composition was slightly higher 1 year after RYGB (17).

While the results of Wolfe et al. (3) add to the literature on metabolic adaptation, we still need to explain how much of the variability in long-term metabolic adaptation is due to differences between cohorts, interventions, or the mathematical models predicting expected EE. **○**

Published 2018. This article is a U.S. Government work and is in the public domain in the USA.

References

1. Elia M. Organ and tissue contribution to metabolic rate. In: Kinney JM, Tucker HN, eds. *Energy Metabolism: Tissue Determinants and Cellular Corollaries*. New York, NY: Raven Press; 1992:61-79.
2. Hall KD. Predicting metabolic adaptation, body weight change, and energy intake in humans. *Am J Physiol Endocrinol Metab* 2010;298:E449-E466.
3. Wolfe BM, Schoeller DA, McCrady-Spitzer SK, Thomas DM, Sorenson CE, Levine JA. Resting metabolic rate, total daily energy expenditure, and metabolic adaptation 6 months and 24 months after bariatric surgery. *Obesity (Silver Spring)* 2018;26:862-868.
4. Knuth ND, Johannsen DL, Tamboli RA, et al. Metabolic adaptation following massive weight loss is related to the degree of energy imbalance and changes in circulating leptin. *Obesity (Silver Spring)* 2014;22:2563-2569.
5. Fothergill E, Guo J, Howard L, et al. Persistent metabolic adaptation 6 years after “The Biggest Loser” competition. *Obesity (Silver Spring)* 2016;24:1612-1619.
6. Pontzer H. Constrained total energy expenditure and the evolutionary biology of energy balance. *Exerc Sport Sci Rev* 2015;43:110-116.
7. Butte NF, Brandt ML, Wong WW, et al. Energetic adaptations persist after bariatric surgery in severely obese adolescents. *Obesity (Silver Spring)* 2015;23:591-601.
8. Tam CS, Redman LM, Greenway F, LeBlanc KA, Haussmann MG, Ravussin E. Energy metabolic adaptation and cardiometabolic improvements one year after gastric bypass, sleeve gastrectomy, and gastric band. *J Clin Endocrinol Metab* 2016;101:3755-3764.
9. Tam CS, Rigas G, Heilbronn LK, Matisan T, Probst Y, Talbot M. Energy adaptations persist 2 years after sleeve gastrectomy and gastric bypass. *Obes Surg* 2016;26:459-463.
10. van Gemert WG, Westerterp KR, van Acker BA, et al. Energy, substrate and protein metabolism in morbid obesity before, during and after massive weight loss. *Int J Obes Relat Metab Disord* 2000;24:711-718.
11. Westerterp KR, Saris WH, Soeters PB, ten Hoor F. Determinants of weight loss after vertical banded gastroplasty. *Int J Obes* 1991;15:529-534.
12. Keys A. *The Biology of Human Starvation*. Minneapolis, MN: University of Minnesota Press; 1950.
13. Rosenbaum M, Hirsch J, Gallagher DA, Leibel RL. Long-term persistence of adaptive thermogenesis in subjects who have maintained a reduced body weight. *Am J Clin Nutr* 2008;88:906-912.

Laboratory of Biological Modeling, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland, USA. Correspondence: Kevin D. Hall (kevinh@niddk.nih.gov)

See accompanying article, pg. 862.

Funding agencies: This research was supported by the Intramural Research Program of the National Institute of Diabetes and Digestive and Kidney Diseases.

Disclosure: Dr. Hall holds a patent on a method of personalized dynamic feedback control of body weight (US Patent No. 9,569,483; assigned to the National Institutes of Health).

Received: 16 March 2018; **Accepted:** 16 March 2018; **Published online** 10 April 2018. doi:10.1002/oby.22189

14. Amatruda JM, Statt MC, Welle SL. Total and resting energy expenditure in obese women reduced to ideal body weight. *J Clin Invest* 1993;92:1236-1242.
15. Das SK, Roberts SB, McCrory MA, et al. Long-term changes in energy expenditure and body composition after massive weight loss induced by gastric bypass surgery. *Am J Clin Nutr* 2003;78:22-30.
16. Weinsier RL, Nagy TR, Hunter GR, Darnell BE, Hensrud DD, Weiss HL. Do adaptive changes in metabolic rate favor weight regain in weight-reduced individuals? An examination of the set-point theory. *Am J Clin Nutr* 2000;72:1088-1094.
17. Wilms B, Ernst B, Thurnheer M, Schmid SM, Spengler CM, Schultes B. Resting energy expenditure after Roux-en Y gastric bypass surgery. *Surg Obes Relat Dis* 2018;14:191-199.